

(19)



Europäisches Patentamt  
European Patent Office  
Office européen des brevets



(11)

**EP 1 382 613 A1**

(12)

**EUROPEAN PATENT APPLICATION**

(43) Date of publication:

**21.01.2004 Bulletin 2004/04**

(51) Int Cl.7: **C07K 14/47**

(21) Application number: **03015519.6**

(22) Date of filing: **09.07.2003**

(84) Designated Contracting States:

**AT BE BG CH CY CZ DE DK EE ES FI FR GB GR  
HU IE IT LI LU MC NL PT RO SE SI SK TR**

Designated Extension States:

**AL LT LV MK**

(30) Priority: **16.07.2002 JP 2002206841**

**19.12.2002 JP 2002367778**

**31.03.2003 JP 2003095955**

(71) Applicant: **Sumitomo Chemical Company,  
Limited**

**Chuo-ku Osaka 541-8550 (JP)**

(72) Inventors:

- **Takahashi, Yasuhiko**  
**Toyonaka-shi, Osaka (JP)**
- **Matsumoto, Yasuo**  
**Toyonaka-shi, Osaka (JP)**
- **Oeda, Kenji**  
**Kyoto-shi, Kyoto (JP)**

(74) Representative: **VOSSIUS & PARTNER**

**Siebertstrasse 4  
81675 München (DE)**

(54) **Novel G protein, polynucleotide encoding the same and utilization thereof**

(57) We identified a novel protein (Gm1) comprising an amino acid sequence part having a high homology with a domain having a high homology with a GTP binding site and a GTPase site conserved among G protein  $\alpha$  subunits and a trimer forming domain conserved among G protein  $\alpha$  subunits. The Gm1 is involved in an signal transduction via a G protein-coupled receptor (GPCR) stimulation. Accordingly, this protein is consid-

ered to be a novel G protein. The Gm1 is expressed intensively in human brain, thymus, testes, spleen, small intestine, uterus and heart. We also established a method for screening for a substance capable of regulating a cellular signal transduction employing a polynucleotide encoding the Gm1.

**EP 1 382 613 A1**

## Description

## BACKGROUND OF THE INVENTION

## 5 Field of the Invention

**[0001]** The present invention relates to a novel G protein having an ability of amplifying a signal transduction generated by a receptor upon binding to a natural ligand such as amine, peptide, hormone, autacoid, neurotransmitter and the like, as well as a polynucleotide encoding the same. The invention also relates to a method for screening for a substance which regulates the cellular signal transduction mediated by this novel G protein, and the like.

## Description of the Background Art

**[0002]** A G protein is an important mediator in a signal transduction. Thus, the G protein serves as a transmitter which transport, into a cell, a stimulation signal received by a G protein-coupled receptor (hereinafter sometimes abbreviated as GPCR) which is a seven transmembrane receptor. A GPCR is expressed in a wide variety of tissues, and this signal transduction system was proven to be involved in regulation of a wide variety of cellular functions such as hormone reception, neurotransmission, cell proliferation and differentiation and the like (GENDAI KAGAKU ZOKAN 34, 61 to 70, 1997).

**[0003]** More specifically, a G protein is a heterotrimer formed from  $\alpha$  subunit which binds to GTP,  $\beta$  subunit and  $\gamma$  subunit. When a GPCR on a cell membrane surface binds to a ligand such as a hormone or neurotransmitter, the GPCR is activated and its signal is transmitted to a G protein. In the G protein to which the signal has been transmitted, a GDP is released from the inactive form in which a GDP was bound once to the  $\alpha$  subunit, and a GTP is then bound instead, whereby converting into an active form.

**[0004]** The active G protein is released from the GPCR while dissociating into a GTP-binding  $\alpha$  subunit and  $\beta\gamma$  subunits. The active G protein promotes or inhibits its target effector such as adenylate cyclase,  $\text{Ca}^{2+}$  channel,  $\text{K}^{+}$  channel, phospholipase  $\text{C}\beta$  and the like, whereby regulating a variety of the cellular functions. A mammalian G protein  $\alpha$  subunit may for example be  $\text{Gi}$ ,  $\text{Go}$ ,  $\text{Gq}$ ,  $\text{Gt}$  and the like. Typically, a G protein  $\alpha$  subunit is classified into any of 4 types, namely, the type which promotes the activity of an adenylate cyclase, the type which inhibits the activity of an adenylate cyclase, the type which promotes the activity of a phospholipase and the type which transmits a signal to a Rho.

**[0005]** A GTP which has been bound to an  $\alpha$  subunit of an active G protein is converted into a GDP by the GTPase effect possessed by the  $\alpha$  subunit, resulting in the recovery of an inactive form ("Signal Dentatsu", p.17-30, Nov. 1, 2001, KYORITSU SHUPPANSHA).

**[0006]** GPCR genes and its gene products and GPCR signal transduction pathway-related genes and its gene products are potential causes for diseases (Spiegel et al., J.Clin.Invest.92: 1119-1125 (1993); MuKusick et al., J.Med.Genet. 30: 1-26 (1993); Lania et al., European J.Endocrinology 145:543-559(2001)). For example, a certain defect in a V2 vasopressin receptor gene as a GPCR has been proven to induce various forms of a nephrogenic diabetes insipidus (Holtzman et al., Hum. Mol. Genet. 2:1201-1204(1993)). In addition, variation in  $\text{G}\alpha$  subunits are observed in a tumor of growth hormone secreting cells in a pituitary gland which secretes a growth hormone, hyperthyroid tumor, ovarian and renal tumors (Meij, JTA (1996), Mol. Cell. Biochem. 157:31-38; Aussel, C. et al., (1988), J.Immunol.140: 215-220). Accordingly, GPCR signal-related gene products are useful as a target of a novel drug, and 50% of currently marketed pharmaceuticals were reported to direct the GPCR as a target (Nature Review Drug Discovery, 1, 7 (2002)).

**[0007]** Generally in screening natural ligands of GPCR, it is important to know which G proteins are to be coupled with the GPCR (Trends in Pharmacological Science, 22, 560-564 (2001)). Accordingly, identification of a novel G protein and a gene encoding the same is useful in treating or diagnosing a disease caused by the abnormality in the cellular signal transduction in which said G protein is involved. In addition, it can be used in the screening for a pharmaceutical which is useful as a remedial, therapeutic or prophylactic agent against a disease caused by the abnormality in a cellular signal transduction. Furthermore, it can be used in the screening for a pharmaceutical which is useful as a remedial, therapeutic or prophylactic agent capable of ameliorating or preventing a symptom by means of the activation or inhibition of the cellular signal transduction.

## SUMMARY OF THE INVENTION

**[0008]** A main object of the present invention is to provide a novel G protein  $\alpha$  subunit and a polynucleotide encoding the same, a substance capable of activating or inhibiting the signal transduction system mediated by a G protein-coupled receptor and this G protein  $\alpha$  subunit, a method for screening for such a substance as well as a screening kit therefor.

[0009] For the purpose of accomplishing the objective described above, we made an effort and finally discovered a human novel protein comprising an amino acid sequence having a high homology with the GTP binding site and the GTPase activation site conserved among G protein  $\alpha$  subunits and an amino acid sequence having a high homology with the heterotrimer forming domain conserved among the G protein  $\alpha$  subunits, and designated this protein as a Gm1 protein. We also discovered a mouse Gm1 protein and a rat Gm1 protein having similar characteristics with regard to the amino acid sequences.

[0010] Moreover, we discovered that in a cell having a Gm1 expression vector, the effector activity of the G protein is elevated.

[0011] Furthermore, we discovered that Gm1 protein is expressed at a high level in human brain, thymus, testes, spleen, small intestine, uterus and heart.

[0012] Based on the findings described above, we believed that the present protein (Gm1 protein) is a novel G protein  $\alpha$  subunit which is a molecule involved in the signal transduction mediated by a GPCR stimulation which functions in human brain, thymus, testes, spleen, small intestine, uterus and heart, thus establishing the present invention.

[0013] Thus, the invention provides the proteins and polynucleotides and the like, which are listed in the following respective paragraphs:

1. A protein comprising any amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence represented by SEQ ID No:1;
- (b) an amino acid sequence of a protein involved in a G protein-coupled receptor mediated signal transduction, said protein consists of an amino acid sequence having a homology of 85% or more with the amino acid sequence represented by SEQ ID No:1;
- (c) the amino acid sequence represented by SEQ ID No:25;
- (d) the amino acid sequence represented by SEQ ID No:26;
- (e) an amino acid sequence of a protein involved in a G protein-coupled receptor mediated signal transduction, said protein comprises the amino acid sequence of the amino acid Nos.96 to 126 of SEQ ID No:1;
- (f) an amino acid sequence of a protein involved in a G protein-coupled receptor mediated signal transduction, said protein comprises an amino acid sequence having a homology of 95% or more with the amino acid sequence of the amino acid Nos.96 to 126 of SEQ ID No:1;
- (g) an amino acid sequence of a protein involved in a G protein-coupled receptor mediated signal transduction, said protein comprises at its N-terminal the amino acid sequence of the amino acid Nos.1 to 126 of SEQ ID No:1; and
- (h) an amino acid sequence of a protein involved in a G protein-coupled receptor mediated signal transduction, said protein comprises at its N-terminal an amino acid sequence having a homology of 65% or more with the amino acid sequence of the amino acid Nos.1 to 126 of SEQ ID No:1.

2. A protein (1) or (2):

- (1) the protein consisting of the amino acid sequence represented by SEQ ID No:1;
- (2) a protein involved in a G protein-coupled receptor mediated signal transduction which consists of an amino acid sequence having a homology of 85% or more with the amino acid sequence represented by SEQ ID No:1.

3. The protein consisting of the amino acid sequence represented by SEQ ID No:25.

4. The protein consisting of the amino acid sequence represented by SEQ ID No:26.

5. A protein (3) or (4):

- (3) a protein involved in a G protein-coupled receptor mediated signal transduction which comprises the amino acid sequence of the amino acid Nos.96 to 126 of SEQ ID No:1;
- (4) a protein involved in a G protein-coupled receptor mediated signal transduction which comprises an amino acid sequence having a homology of 95% or more with the amino acid sequence of the amino acid Nos.96 to 126 of SEQ ID No:1.

6. A protein (5) or (6):

- (5) a protein involved in a G protein-coupled receptor mediated signal transduction which comprises at its N-terminal the amino acid sequence of the amino acid Nos.1 to 126 of SEQ ID No:1;
- (6) a protein involved in a G protein-coupled receptor mediated signal transduction which comprises at its N-terminal an amino acid sequence having a homology of 65% or more with the amino acid sequence of the

amino acid Nos.1 to 126 of SEQ ID No:1.

7. A polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) a nucleotide sequence encoding the amino acid sequence represented by SEQ ID No:1;
- (b) a nucleotide sequence encoding an amino acid sequence of a protein involved in a G protein-coupled receptor mediated signal transduction, said protein consists of an amino acid sequence having a homology of 85% or more with the amino acid sequence represented by SEQ ID No:1;
- (c) a nucleotide sequence encoding the amino acid sequence represented by SEQ ID No:25;
- (d) a nucleotide sequence encoding the amino acid sequence represented by SEQ ID No:26;
- (e) a nucleotide sequence encoding an amino acid sequence of a protein involved in a G protein-coupled receptor mediated signal transduction, said protein comprises the amino acid sequence of the amino acid Nos.96 to 126 of SEQ ID No:1;
- (f) a nucleotide sequence encoding an amino acid sequence of a protein involved in a G protein-coupled receptor mediated signal transduction, said protein comprises an amino acid sequence having a homology of 95% or more with the amino acid sequence of the amino acid Nos.96 to 126 of SEQ ID No:1;
- (g) a nucleotide sequence encoding an amino acid sequence of a protein involved in a G protein-coupled receptor mediated signal transduction, said protein comprises at its N-terminal the amino acid sequence of the amino acid Nos.1 to 126 of SEQ ID No:1;
- (h) a nucleotide sequence encoding an amino acid sequence of a protein involved in a G protein-coupled receptor mediated signal transduction, said protein comprises at its N-terminal an amino acid sequence having a homology of 65% or more with the amino acid sequence of the amino acid Nos.1 to 126 of SEQ ID No:1;
- (i) the nucleotide sequence represented by SEQ ID No:2;
- (j) a nucleotide sequence of a polynucleotide encoding a protein involved in a G protein-coupled receptor mediated signal transduction, said polynucleotide consists of a nucleotide sequence having a homology of 85% or more with the polynucleotide consisting of the nucleotide sequence represented by SEQ ID No:2;
- (k) the nucleotide sequence represented by SEQ ID No:27;
- (l) the nucleotide sequence represented by SEQ ID No:28;
- (m) a nucleotide sequence of a polynucleotide encoding a protein involved in a G protein-coupled receptor mediated signal transduction, said polynucleotide comprises the nucleotide sequence of the nucleotide Nos.289 to 378 of SEQ ID No:2;
- (n) a nucleotide sequence of a polynucleotide encoding a protein involved in a G protein-coupled receptor mediated signal transduction, said polynucleotide comprises a nucleotide sequence having a homology of 90% or more with the polynucleotide consisting of the nucleotide sequence of the nucleotide Nos.289 to 378 of SEQ ID No:2;
- (o) a nucleotide sequence of a polynucleotide encoding a protein involved in a G protein-coupled receptor mediated signal transduction, said polynucleotide comprises at its 5' terminal the nucleotide sequence of the nucleotide Nos.1 to 378 of SEQ ID No:2; and
- (p) a nucleotide sequence of a polynucleotide encoding a protein involved in a G protein-coupled receptor mediated signal transduction, said polynucleotide comprises at its 5' terminal a nucleotide sequence having a homology of 70% or more with the polynucleotide consisting of the nucleotide sequence of the nucleotide Nos.1 to 378 of SEQ ID No:2.

8. A polynucleotide of (7) or (8):

- (7) the polynucleotide consisting of the nucleotide sequence represented by SEQ ID No:2;
- (8) a polynucleotide encoding a protein involved in a G protein-coupled receptor mediated signal transduction, said polynucleotide consists of a nucleotide sequence having a homology of 85% or more with the polynucleotide consisting of the nucleotide sequence represented by SEQ ID No:2.

9. The polynucleotide consisting of the nucleotide sequence represented by SEQ ID No:27.

10. The polynucleotide consisting of the nucleotide sequence represented by SEQ ID No:28.

11. A polynucleotide of (9) or (10):

- (9) a polynucleotide encoding a protein involved in a G protein-coupled receptor mediated signal transduction, said polynucleotide comprises the nucleotide sequence of the nucleotide Nos.289 to 378 of SEQ ID No:2;
- (10) a polynucleotide encoding a protein involved in a G protein-coupled receptor mediated signal transduction, said polynucleotide comprises a nucleotide sequence having a homology of 90% or more with the polynucleotide

otide consisting of the nucleotide sequence of the nucleotide Nos.289 to 378 of SEQ ID No:2.

12. A polynucleotide of (11) or (12):

(11) a polynucleotide encoding a protein involved in a G protein-coupled receptor mediated signal transduction, said polynucleotide comprises at its 5' terminal the nucleotide sequence of the nucleotide Nos.1 to 378 of SEQ ID No:2;

(12) a polynucleotide encoding a protein involved in a G protein-coupled receptor mediated signal transduction, said polynucleotide comprises at its 5' terminal a nucleotide sequence having a homology of 70% or more with the polynucleotide consisting of the nucleotide sequence of the nucleotide Nos.1 to 378 of SEQ ID No:2.

13. A recombinant vector containing a polynucleotide according to the above 7.

14. A method for producing a recombinant vector comprising a step for integrating a polynucleotide according to the above 7 into a vector capable of being replicated in a host cell.

15. A transformant having a recombinant vector according to the above 13.

16. A method for producing a transformant comprising a step for transducing a recombinant vector according to the above 13 into a host cell.

17. A method for producing a G protein  $\alpha$ -subunit comprising steps for culturing a transformant having a recombinant vector containing a polynucleotide according to the above 7 and recovering from the culture a protein resulting from the polynucleotide according to the above 7.

18. An antisense polynucleotide consisting of a polynucleotide of (13) or (14):

(13) a polynucleotide which inhibits the expression of a protein according to the above 1 which comprises a nucleotide sequence complementary to at least 15 contiguous nucleotides in the nucleotide sequence represented by SEQ ID No:2;

(14) a polynucleotide which inhibits the expression of a protein according to the above 1 which hybridizes under an intracellular condition with a polynucleotide consisting of at least 15 contiguous nucleotides in the nucleotide sequence represented by SEQ ID No:2.

19. A ribozyme (15) or (16):

(15) a ribozyme having an ability of cleaving a polynucleotide according to the above 7 which comprises two polynucleotide regions complementary to two regions respectively consisting of at least 9 contiguous nucleotides which are two regions in the nucleotide sequence represented by SEQ ID No:2;

(16) a ribozyme having an ability of cleaving a polynucleotide according to the above 7 which comprises two polynucleotide regions which hybridizes under an intracellular condition with two regions respectively consisting of at least 9 contiguous nucleotides which are two regions in the nucleotide sequence represented by SEQ ID No:2.

20. An antibody which recognizes a protein according to the above 1 specifically.

21. An agent for regulating a G protein-coupled receptor mediated signal transduction containing as an active ingredient a protein according to the above 1.

22. A therapeutic or prophylactic agent against a disease caused by a G protein-coupled receptor mediated signal transduction abnormality, wherein an active ingredient of the agent is a protein according to the above 1.

23. An agent for regulating a G protein-coupled receptor mediated signal transduction containing as an active ingredient a polynucleotide according to the above 7.

24. A therapeutic or prophylactic agent against a disease caused by a G protein-coupled receptor mediated signal transduction abnormality, wherein an active ingredient of the agent is a polynucleotide according to the above 7.

25. An agent for regulating a G protein-coupled receptor mediated signal transduction containing as an active ingredient an antisense polynucleotide according to the above 18.

26. A therapeutic or prophylactic agent against a disease caused by a G protein-coupled receptor mediated signal transduction abnormality, wherein an active ingredient of the agent is an antisense polynucleotide according to the above 18.

27. An agent for regulating a G protein-coupled receptor mediated signal transduction containing as an active ingredient a ribozyme according to the above 19.

28. A therapeutic or prophylactic agent against a disease caused by a G protein-coupled receptor mediated signal transduction abnormality, wherein an active ingredient of the agent is a ribozyme according to the above 19.

29. An agent for regulating a G protein-coupled receptor mediated signal transduction containing as an active

ingredient an antibody according to the above 20.

30. A therapeutic or prophylactic agent against a disease caused by a G protein-coupled receptor mediated signal transduction abnormality, wherein an active ingredient of the agent is an antibody according to the above 20.

31. An oligonucleotide (17) or (18):

- (17) an oligonucleotide capable of recognizing a polynucleotide represented by SEQ ID NO:2 specifically which consists of at least 17 contiguous nucleotides in the nucleotide sequence represented by SEQ ID No:2;
- (18) an oligonucleotide capable of recognizing a polynucleotide represented by SEQ ID NO:2 specifically which has a homology of 80% or more with at least 17 contiguous nucleotides in the nucleotide sequence represented by SEQ ID No:2.

32. An oligonucleotide according to the above 31 which is used as a probe or a primer.

33. A method for screening for a substance capable of regulating a signal transduction mediated by a G protein-coupled receptor and a G protein comprising:

- (a) a step for bringing a test substance into contact with a test cell having a recombinant vector according to the above 13 and a recombinant vector containing a DNA encoding a G protein-coupled receptor protein;
- (b) a step for measuring the G protein effector activity or the index value correlating therewith in the test cell; and
- (c) a step for comparing this effector activity or the index value correlating therewith with the effector activity or the index value correlating therewith in the test cell which has not been brought into contact with the test substance, whereby selecting a test substance capable of altering the effector activity or the index value correlating therewith in the test cell.

34. A method for screening for a substance capable of regulating a signal transduction mediated by a G protein-coupled receptor and a G protein comprising:

- (a) a step for bringing a test substance into contact with a test cell having a recombinant vector according to the above 13 and a recombinant vector containing a DNA encoding a G protein-coupled receptor protein;
- (b) a step for measuring the G protein effector activity or the index value correlating therewith in the test cell; and
- (c) a step for comparing this effector activity with the effector activity or the index value correlating therewith when the said test substance has been brought into contact with a control cell having no recombinant vector according to the above 13 but having a recombinant vector containing a DNA encoding a G protein-coupled receptor protein, whereby selecting a test substance causing a difference in the effector activity or the index value correlating therewith between the test cell and the control cell.

35. A method for screening for a substance capable of regulating a signal transduction mediated by a G protein-coupled receptor and a G protein comprising:

- (a) a step for bringing a test substance into contact with a test cell having a recombinant vector according to the above 13 and a recombinant vector containing a DNA encoding a G protein-coupled receptor protein;
- (b) a step for measuring the G protein effector activity or the index value correlating therewith in the test cell; and
- (c) a step for comparing this effector activity or the index value correlating therewith with the effector activity or the index value correlating therewith when the said test substance has been brought into contact with a control cell having no recombinant vector containing a DNA encoding a G protein-coupled receptor protein but having a recombinant vector according to the above 13, whereby selecting a test substance causing a difference in the effector activity or the index value correlating therewith between the test cell and the control cell.

36. A method for screening for a substance capable of regulating a signal transduction mediated by a G protein-coupled receptor and a G protein comprising:

- (a) a step for bringing a test substance and a G protein-coupled receptor ligand into contact with a test cell having a recombinant vector according to the above 13 and a recombinant vector containing a DNA encoding a G protein-coupled receptor protein;
- (b) a step for measuring the G protein effector activity or the index value correlating therewith in the test cell; and
- (c) a step for comparing this effector activity or the index value correlating therewith with the effector activity or the index value correlating therewith in the test cell which has not been brought into contact with the test substance but has been brought into contact with the ligand, whereby selecting a test substance capable of

altering the effector activity or the index value correlating therewith in the test cell.

37. A method for screening for a substance capable of regulating a signal transduction mediated by a G protein-coupled receptor and a G protein comprising:

- (a) a step for bringing a test substance and a G protein-coupled receptor ligand into contact with a test cell having a recombinant vector according to the above 13 and a recombinant vector containing a DNA encoding a G protein-coupled receptor protein;
- (b) a step for measuring the G protein effector activity or the index value correlating therewith in the test cell;
- (c) a step for comparing this effector activity with the effector activity in the test cell which has not been brought into contact with the test substance but has been brought into contact with the ligand, whereby investigating the change in the effector activity in the test cell; and
- (d) a step for comparing the rate of change in this effector activity or the index value correlating therewith with the rate of change in the effector activity or the index value correlating therewith when the said test substance and said ligand has been brought into contact with a control cell having no recombinant vector containing a DNA encoding a G protein-coupled receptor protein but having a recombinant vector according to the above 13, whereby selecting a test substance causing a difference in the rate of change in the effector activity or the index value correlating therewith between the test cell and the control cell.

38. A method for screening for a substance capable of regulating a signal transduction mediated by a G protein-coupled receptor and a G protein comprising:

- (a) a step for bringing a test substance into contact with a cell membrane fraction of a cell having a recombinant vector according to the above 13 and a cell membrane fraction of a cell having a recombinant vector containing a DNA encoding a GPCR, or a cell membrane fraction of a cell having the recombinant vector according to the above 13 and the recombinant vector containing the DNA encoding the GPCR;
- (b) a step for assaying the level of the binding of GTP to the cell membrane fraction; and
- (c) a step for comparing the assayed level of this GTP binding with the assayed level of the GTP binding to the cell membrane fraction which has not been brought into contact with the test substance, whereby selecting a test substance capable of altering the assayed level of the GTP binding to the cell membrane fraction.

39. A method for screening for a substance capable of regulating a signal transduction mediated by a G protein-coupled receptor and a G protein comprising:

- (a) a step for bringing a test substance and a G protein-coupled receptor ligand into contact with a cell membrane fraction of a cell having a recombinant vector according to the above 13 and a cell membrane fraction of a cell having a recombinant vector containing a DNA encoding a GPCR, or a cell membrane fraction of a cell having the recombinant vector according to the above 13 and the recombinant vector containing the DNA encoding the GPCR;
- (b) a step for assaying the level of the binding of GTP to the cell membrane fraction; and
- (c) a step for comparing the assayed level of this GTP binding with the assayed level of the GTP binding in the cell membrane fraction which has not been brought into contact with the test substance but has been brought into contact with said ligand, whereby selecting a test substance capable of altering the assayed level of the GTP binding to the cell membrane fraction.

40. A method for screening for a substance capable of regulating a signal transduction mediated by a G protein-coupled receptor and a G protein comprising:

- (a) a step for bringing a test substance into contact with a test cell capable of expressing a protein according to the above 1;
- (b) a step for measuring the expression level of the protein according to the above 1 in the test cell; and
- (c) a step for comparing this expression level with the expression level of said protein in the test cell which has not been brought into contact with the test substance, whereby selecting a test substance capable of altering the expression level of said protein in the test cell.

41. A substance capable of regulating a signal transduction mediated by a G protein-coupled receptor and a G protein, said substance is obtained by a screening method according to any of the above 33 to 40.

42. An agent for regulating a signal transduction mediated by a G protein-coupled receptor and a G protein, said agent contains as an active ingredient a substance according to the above 41.

43. A therapeutic or prophylactic agent against a disease caused by the abnormality in a G protein-coupled receptor and a G protein-mediated signal transduction containing as an active ingredient a substance according to the above 41.

44. A kit for screening for a substance capable of regulating a signal transduction mediated by a G protein-coupled receptor and a protein according to the above 1, which comprises a test cell having a recombinant vector containing a polynucleotide encoding a protein according to the above 1 and a reagent for measuring the G protein effector activity or an index value correlating therewith.

45. A screening kit according to the above 44 wherein the test cell further has a recombinant vector having a DNA encoding a G protein-coupled receptor.

46. A screening kit according to the above 44 further containing a G protein-coupled receptor ligand.

47. A screening kit according to the above 44 further containing a control cell having a recombinant vector having a DNA encoding a G protein-coupled receptor.

48. A screening kit according to the above 44 further containing a control cell having a recombinant vector containing a polynucleotide encoding a protein according to the above 1.

49. A kit for screening for a substance capable of regulating a signal transduction mediated by a G protein-coupled receptor and a protein according to the above 1, which comprises:

a cell having a recombinant vector containing a polynucleotide encoding a protein according to the above 1;  
and,  
a GTP analogue which can bind to the protein according to the above 1 but can not be cleaved by a GTPase.

50. A screening kit according to the above 49 wherein the cell further has a recombinant vector having a DNA encoding a G protein-coupled receptor.

51. A kit for screening for a substance capable of regulating a signal transduction mediated by a G protein-coupled receptor and a protein according to the above 1, which comprises:

a cell having a recombinant vector containing a polynucleotide encoding a protein according to the above 1;  
a cell having a recombinant vector having a DNA encoding the G protein-coupled receptor; and,  
a GTP analogue which can bind to the protein according to the above 1 but can not be cleaved by a GTPase.

52. A screening kit according to the above 49 or 51 further containing a G protein-coupled receptor ligand.

53. A kit for screening for a substance capable of regulating a signal transduction mediated by a G protein-coupled receptor and a protein according to the above 1, which comprises:

a cell capable of expressing a protein according to the above 1; and  
an oligonucleotide according to the above 31 or an antibody according to the above 20.

## BRIEF DESCRIPTION OF THE DRAWINGS

[0014]

Figure 1 is a drawing substitute showing the expression profile of an inventive protein in human tissues;

Figure 2 is a schematic view indicating the dopamine D1 receptor antagonistic effects of various test substance (Example 17);

Figure 3 is a schematic view indicating the dopamine D1 receptor agonistic effects of various test substance (Example 18);

Figure 4 is a schematic view indicating the adenosine A2a receptor antagonistic effects of DMPX (Example 21);

Figure 5 is a schematic view indicating the adenosine A2a receptor agonistic effects of CGS-21680 (Example 22);

Figure 6 is a schematic view indicating that dopamine caused a signal transduction mediated by the dopamine D1 receptor and the Gm1 (Example 24); and

Figure 7 is a schematic view indicating that adenosine caused a signal transduction mediated by the adenosine A2a receptor and the Gm1 (Example 24).



## DESCRIPTION OF THE PREFERRED EMBODIMENTS

## &lt;Inventive proteins&gt;

5 **[0015]** An inventive protein is a protein comprising any amino acid sequence selected from the group consisting of:

- (a) The amino acid sequence represented by SEQ ID No:1;
- (b) An amino acid sequence of a protein involved in a G protein-coupled receptor mediated signal transduction which consists of an amino acid sequence having a homology of 85% or more with the amino acid sequence represented by SEQ ID No: 1;
- (c) The amino acid sequence represented by SEQ ID No:25;
- (d) The amino acid sequence represented by SEQ ID No:26;
- (e) An amino acid sequence of a protein involved in a G protein-coupled receptor mediated signal transduction which comprises the amino acid sequence of the amino acid Nos.96 to 126 of SEQ ID No:1;
- 10 (f) An amino acid sequence of a protein involved in a G protein-coupled receptor mediated signal transduction which comprises an amino acid sequence having a homology of 95% or more with the amino acid sequence of the amino acid Nos.96 to 126 of SEQ ID No:1;
- (g) An amino acid sequence of a protein involved in a G protein-coupled receptor mediated signal transduction which comprises at its N-terminal the amino acid sequence of the amino acid Nos.1 to 126 of SEQ ID No:1; and
- 15 (h) An amino acid sequence of a protein involved in a G protein-coupled receptor mediated signal transduction which comprises at its N-terminal an amino acid sequence having a homology of 65% or more with the amino acid sequence of the amino acid Nos.1 to 126 of SEQ ID No:1;
- 20

**[0016]** The inventive protein includes not only said protein but also its salt or derivative as long as its biological functions are not affected. A derivative is not limited particularly and may for example be one whose C terminal or other carboxyl group is converted into an amide, ester and the like, or one whose N terminal or other amino group is protected for example by a formyl group or acyl group. As a salt, an acid addition salt is exemplified preferably. An acid addition salt may for example be a salt with an inorganic acid such as hydrochloric acid, phosphoric acid, sulfuric acid and the like, or a salt with an organic acid such as formic acid, acetic acid, propionic acid and the like.

## &lt;First protein&gt;

**[0017]** The first protein according to the invention is a protein (1) or (2) shown below.

- 35 (1) The protein consisting of the amino acid sequence represented by SEQ ID No:1.
- (2) A protein involved in a G protein-coupled receptor mediated signal transduction which consists of an amino acid sequence having a homology of 85% or more with the amino acid sequence represented by SEQ ID No:1.

**[0018]** The amino acid sequence represented by SEQ ID No:1 comprises an amino acid sequence part having a high homology with the GTP binding site and the GTPase activation site conserved among G protein  $\alpha$  subunits. Such parts are the regions of the amino acid Nos.126 to 133, 287 to 292, 353 to 359, 428 to 435 in the amino acid sequence represented by SEQ ID No:1. Any of these amino acids is in agreement with the GTP binding site and the GTPase activation site of Gs and Golf which has already been identified as G protein  $\alpha$  subunits (NATURE, 117-127, 1991, vol. 349).

45 **[0019]** Furthermore, the amino acid also comprises a sequence which is identical to the characteristic sequence conserved highly in Gs and Golf protein belonging especially to the Gs family among the G protein  $\alpha$  subunits (amino acid Nos.119 to 126 in SEQ ID No:1), and can also form an  $\alpha$  helix structure conserved among the G protein  $\alpha$  subunits.

**[0020]** Based on these findings, the protein comprising the amino acid sequence represented by SEQ ID No:1 is considered to be a G protein  $\alpha$  subunit.

50 **[0021]** The fact that a protein (2) functions as a molecule involved in the intracellular signal transduction by a GPCR stimulation can be verified by means of a screening method according to the invention discussed below.

**[0022]** The amino acid sequence of a protein (2) preferably has a homology of 90% or more, especially 95% or more with the amino acid sequence represented by SEQ ID No:1.

**[0023]** An index indicating which and how many amino acid residues in a protein (2) can be substituted, deleted or added without losing any biological functions can be identified for example by a GTP binding level assay described below. A variation causing no loss of the biological functions can be conducted for example in a part having a low homology with the amino acid sequence of any of various G protein  $\alpha$  subunits which have already been identified.

**[0024]** In the case for example of an amino acid substitution, the amino acid can be substituted by an amino acid

having the characteristics similar to those of the former amino acid in terms of polarity, electric charge, solubility, hydrophilicity/hydrophobicity, polarity and the like, in view of the maintenance of the protein structure. In this context, glycine, alanine, valine, leucine, isoleucine and proline are classified into non-polar amino acids; serine, threonine, cysteine, methionine, asparagine and glutamine are classified into polar amino acids; phenylalanine, tyrosine and triptophane are classified into aromatic side chain-carrying amino acids; lysine, arginine and histidine are classified into basic amino acids; aspartic acid and glutamic acid are classified into acidic amino acids. Accordingly, the substitution may be conducted using an amino acid selected from the same group.

**[0025]** A protein (2) also includes proteins derived from other species corresponding to the human protein. Such an other species-derived corresponding protein can be deduced from a nucleotide sequence identified by means for example of a screening of a gene library of other species using a full length inventive polynucleotide or a part thereof as well as a 5'-RACE. Otherwise, a deductive identification is possible also from a corresponding gene of other species screened by an NCBI Blast search described below.

**[0026]** A protein (2) may for example be the protein consisting of the amino acid sequence represented by SEQ ID No:25 and the protein consisting of the amino acid sequence represented by SEQ ID No:26.

<Second protein>

**[0027]** The second protein according to the invention is a protein (3) or (4) shown below.

(3) A protein involved in a G protein-coupled receptor mediated signal transduction which comprises the amino acid sequence of the amino acid Nos.96 to 126 of SEQ ID No:1.

(4) A protein involved in a G protein-coupled receptor mediated signal transduction which comprises an amino acid sequence having a homology of 95% or more with the amino acid sequence of the amino acid Nos.96 to 126 of SEQ ID No:1.

**[0028]** A protein (4) preferably has a homology of 96% or more, especially 97% or more with the amino acid Nos.96 to 126 in the amino acid sequence represented by SEQ ID No:1.

**[0029]** For the purpose of functioning as a molecule involved in an intracellular signal transduction by a GPCR stimulation, for example, as a G protein  $\alpha$  subunit, each of the protein (3) and (4) usually has an amino acid sequence of the amino acid Nos.75 to 133, 287 to 292, 353 to 359 and 428 to 435 in the amino acid sequence represented by SEQ ID No:1, or preferably has an amino acid sequence having a homology usually of 80% or more, especially 90% or more with the amino acid sequence of these regions. The total amino acid number is usually about 320 to 489, preferably about 350 to 460.

**[0030]** An index indicating which and how many amino acid residues in a protein (4) can be substituted, deleted or added without losing any biological functions of the protein (3) can be identified for example by a GTP binding level assay described below. A variation causing no loss of the biological functions can be conducted for example in a part having a low homology with the amino acid sequence of any of various G protein  $\alpha$  subunits which have already been identified. Also similarly to the first protein described above, the substitution of a base can be conducted so that the amino acid obtained after a translation can possess the characteristics analogous to those of the amino acid before the substitution, with regard to polarity, electric charge, solubility, hydrophilicity/hydrophobicity, polarity and the like.

<Third protein>

**[0031]** The third protein of the invention is a protein (5) or (6) shown below.

(5) A protein involved in a G protein-coupled receptor mediated signal transduction which comprises at its N-terminal of the amino acid sequence of the amino acid Nos.1 to 126 of SEQ ID No:1.

(6) A protein involved in a G protein-coupled receptor mediated signal transduction which comprises at its N-terminal of an amino acid sequence having a homology of 65% or more with the amino acid sequence of the amino acid Nos.1 to 126 of SEQ ID No:1.

**[0032]** In a protein (6), the amino acid sequence part corresponding to the 126 amino acid sequence at the N terminal of the protein (5) (hereinafter referred to as a "specific N terminal amino acid sequence") preferably has a homology of 70% or more, especially 75% or more with the specific N terminal amino acid sequence of the protein (5).

**[0033]** For the purpose of functioning as a molecule involved in an intracellular signal transduction by a GPCR stimulation, for example, as a G protein  $\alpha$  subunit, each of the protein (5) and (6) usually has an amino acid sequence of the amino acid Nos.75 to 133, 287 to 292, 353 to 359 and 428 to 435 in the amino acid sequence represented by SEQ ID No:1, or preferably has an amino acid sequence having a homology usually of 80% or more, especially 90% or

more with these regions. The total amino acid number is usually about 320 to 480, preferably about 350 to 460.

[0034] An index indicating which and how many amino acid residues in a protein (6) can be substituted, deleted or added without losing any biological functions of the protein (5) can be identified for example by a GTP binding level assay described below. A variation causing no loss of the biological functions can be conducted for example in a part having a low homology with the amino acid sequence of any of various G protein  $\alpha$  subunits which have already been identified. Also similarly to the first protein described above, the substitution of a base can be conducted so that the amino acid obtained after a translation can possess the characteristics analogous to those of the amino acid before the substitution, with regard to polarity, electric charge, solubility, hydrophilicity/hydrophobicity, polarity and the like.

[0035] There is no known G protein  $\alpha$  subunit having the amino acid sequence of the amino acid Nos.1, to 126 of the amino acid sequence represented by SEQ ID No:1 or a sequence analogous thereto.

#### <Inventive protein production method>

[0036] A protein of the invention can be produced by i) a method for separating a membrane fraction containing said protein from a cell or tissue of a human or other animal species followed by a known protein purification process, ii) a method employing a transformant of the invention described below or iii) a known chemical synthesis of a protein and the like.

[0037] In a method i), a cell or tissue of a mammalian animal including a human can be employed without limitation. It is particularly preferred to use a human cell or tissue, especially, a human brain-, uterus- or heart-derived cell or tissue.

[0038] A membrane fraction containing a protein of the invention can be obtained by suspending a cell or tissue for example in a HE/PI buffer (20mM Hepes, 2mM EDTA, 1 x Proteinase inhibitor cocktail (Nacalaitesque)), pulverizing or lysing the suspension by means for example of an ultrasonic treatment, homogenization, passage through a needle of about 26 gauge, centrifuging the resultant pulverization or lysis solution at about 100 to 150 xG for 5 to 10 minutes, centrifuging the resultant supernatant at about 18,000 to 20,000 G for 20 to 30 minutes, and then recovering the pellets.

[0039] The fact that the resultant cell membrane fraction contains a protein of the invention can be verified for example by a Western blotting using an antibody of the invention as described below.

[0040] A known protein purification method may for example be any of various chromatographic procedure such as an ion exchange, gel filtration, affinity chromatography.

As a method iii), a method described for example in "The Peptide", Academic Press, New York (1966) or a method employing a commercial protein synthesis resin may be exemplified.

[0041] It is also possible to obtain a protein (2) from a transformant having a variant DNA which is formed by imparting a DNA encoding the protein (1) with a variation using a known method such as one described in Molecular Cloning: A Laboratory Manual, 2nd edition, Vol.1 to 3, Cold Spring Harbor Laboratory Press(1989), Methods in Enzymology p448 (1989), PCR A Practical Approach IRL Press p200(1991) and the like, for example, a site-specific mutation introduction, a PCR employing a variation primer. This may analogously be applied to a method for obtaining the protein (3) from a protein (4) and a method for obtaining the protein (6) from a protein (5).

#### <Application of inventive proteins>

[0042] A protein of the invention can preferably be employed as a regulator of a signal transduction mediated by a GPCR stimulation. Specifically, it can preferably be employed for treating or preventing a disease associated with an intracellular signal transduction relating to the defect, reduced expression level or reduced function of a protein of the invention.

#### <Inventive polynucleotide>

[0043] A polynucleotide according to the invention is a polynucleotide encoding a protein of the invention described above, and comprises a nucleotide sequence selected from the group consisting of:

- (a) A nucleotide sequence encoding the amino acid sequence represented by SEQ ID No:1;
- (b) A nucleotide sequence encoding an amino acid sequence of a protein involved in a G protein-coupled receptor mediated signal transduction which consists of an amino acid sequence having a homology of 85% or more with the amino acid sequence represented by SEQ ID No:1;
- (c) A nucleotide sequence encoding the amino acid sequence represented by SEQ ID No:25;
- (d) A nucleotide sequence encoding the amino acid sequence represented by SEQ ID No:26;
- (e) A nucleotide sequence encoding an amino acid sequence of a protein involved in a G protein-coupled receptor mediated signal transduction which comprises the amino acid sequence of the amino acid Nos.96 to 126 of SEQ ID No:1;

(f) A nucleotide sequence encoding an amino acid sequence of a protein involved in a G protein-coupled receptor mediated signal transduction which comprises an amino acid sequence having a homology of 95% or more with the amino acid sequence of the amino acid Nos.96 to 126 of SEQ ID No: 1;

(g) A nucleotide sequence encoding an amino acid sequence of a protein involved in a G protein-coupled receptor mediated signal transduction which comprises at its N-terminal the amino acid sequence of the amino acid Nos. 1 to 126 of SEQ ID No:1;

(h) A nucleotide sequence encoding an amino acid sequence of a protein involved in a G protein-coupled receptor mediated signal transduction which comprises at its N-terminal an amino acid sequence having a homology of 65% or more with the amino acid sequence of the amino acid Nos.1 to 126 of SEQ ID No:1;

(i) The nucleotide sequence represented by SEQ ID No:2;

(j) A nucleotide sequence encoding a protein involved in a G protein-coupled receptor mediated signal transduction which is a nucleotide sequence having a homology of 85% or more with the polynucleotide consisting of the nucleotide sequence represented by SEQ ID No:2;

(k) The nucleotide sequence represented by SEQ ID No:27;

(l) The nucleotide sequence represented by SEQ ID No:28;

(m) A nucleotide sequence encoding a protein involved in a G protein-coupled receptor mediated signal transduction which comprises the nucleotide sequence of the nucleotide Nos.289 to 378 of SEQ ID No:2;

(n) A nucleotide sequence encoding a protein involved in a G protein-coupled receptor mediated signal transduction which comprises a nucleotide sequence having a homology of 90% or more with the polynucleotide consisting of the nucleotide sequence of the nucleotide Nos.289 to 378 of SEQ ID No:2;

(o) A nucleotide sequence encoding a protein involved in a G protein-coupled receptor mediated signal transduction which comprises at its 5' terminal the nucleotide sequence of the nucleotide Nos.1 to 378 of SEQ ID No:2; and

(p) A nucleotide sequence encoding a protein involved in a G protein-coupled receptor mediated signal transduction which comprises at its 5' terminal a nucleotide sequence having a homology of 70% or more with the polynucleotide consisting of the nucleotide sequence of the nucleotide Nos.1 to 378 of SEQ ID No:2.

**[0044]** A polynucleotide of the invention (including an oligonucleotide) includes a polynucleotide comprising a nucleotide sequence described above and a polynucleotide complementary thereto. Unless otherwise specified, a polynucleotide includes the both of a DNA and an RNA. A DNA includes a single-stranded DNA having its nucleotide sequence, and a single-stranded DNA complementary thereto, and a double-stranded DNA. A DNA, unless otherwise specified, includes a cDNA, genome DNA and synthetic DNA. An RNA, unless otherwise specified, includes a total RNA, mRNA, rRNA and synthetic RNA.

**[0045]** An inventive polynucleotide is detailed with referring to the following first, second and third polynucleotide described below.

<First polynucleotide>

**[0046]**

(7) The polynucleotide consisting of the nucleotide sequence represented by SEQ ID No:2.

(8) A polynucleotide encoding a protein involved in a G protein-coupled receptor mediated signal transduction which consists of a nucleotide sequence having a homology of 85% or more with the polynucleotide consisting of the nucleotide sequence represented by SEQ ID No:2.

**[0047]** A polynucleotide (8) preferably has a homology of 87% or more, especially 90% or more with the polynucleotide (7).

**[0048]** A polynucleotide (8) is preferably one which hybridizes under a stringent condition with the polynucleotide (7) consisting of the nucleotide sequence represented by SEQ ID No:2. In the invention, a stringent condition may for example be a condition involving 2 x SSC, 1 x Denhart's solution at about 60°C.

**[0049]** An index indicating which and how many bases in a polynucleotide (8) can be substituted, deleted or added without losing any biological functions of the protein encoded by the polynucleotide (7) can be identified for example by a cAMP assay or a GTP binding level assay described below. A variation causing no loss of the biological functions can be conducted for example in a part having a low homology with the polynucleotide sequence of any of various G protein  $\alpha$  subunits which have already been identified.

**[0050]** Also similarly to the first protein described above, the substitution of a base can be conducted so that the amino acid obtained after a translation can possess the characteristics analogous to those of the amino acid before the substitution, with regard to polarity, electric charge, solubility, hydrophilicity/hydrophobicity, polarity and the like.

**[0051]** When a single amino acid has several translation codons, the base substitution within these translation codons

may also be possible. For example, when alanine has 4 translation codons, namely, GCA, GCC, GCG and GCT, the third base in each codon can be substituted with each other among ATGC.

**[0052]** A polynucleotide (8) includes a polynucleotide of other species corresponding for example to a human polynucleotide. Such a polynucleotide can be screened for using NCBI blast search ([www.ncbi.nlm.nih.gov/BLAST/](http://www.ncbi.nlm.nih.gov/BLAST/)). Typically, the nucleotide sequence containing the nucleotide Nos.289 to 378 in SEQ ID No:2 is subjected to an NCBI blast search whereby searching the nucleotide sequence database of other species and an EST database for a sequence having a high homology. By screening the nucleotide sequences selected by the search for a nucleotide sequence whose region corresponding to the nucleotide Nos.289 to 378 has a homology for example of 90% or more, a corresponding gene of other species can be screened for.

**[0053]** A polynucleotide (8) is preferably one whose nucleotide sequence corresponding to the nucleotide Nos.1 to 222, 400 to 858, 877 to 1056, 1078 to 1281 and 1306 to 1377 in SEQ ID No:2 in the polynucleotide (7) has a homology usually of 75% or more, especially 80% or more with the respective nucleotide sequence of the polynucleotide (7).

**[0054]** A polynucleotide (8) may for example be a polynucleotide consisting of the nucleotide sequence represented by SEQ ID No:27 and a polynucleotide consisting of the nucleotide sequence represented by SEQ ID No:28.

<Second polynucleotide>

**[0055]**

(9) A polynucleotide encoding a protein involved in a G protein-coupled receptor mediated signal transduction which comprises the nucleotide sequence of the nucleotide Nos.289 to 378 of SEQ ID No:2.

(10) A polynucleotide encoding a protein involved in a G protein-coupled receptor mediated signal transduction which comprises a nucleotide sequence having a homology of 90% or more with the polynucleotide consisting of the nucleotide sequence of the nucleotide Nos.289 to 378 of SEQ ID No:2.

**[0056]** In a polynucleotide (10), the nucleotide sequence corresponding to the nucleotide Nos.289 to 378 in SEQ ID No:2 of a polynucleotide (9) has a homology of 93% or more, especially 95% or more with the respective sequence of (9).

**[0057]** A polynucleotide (10) preferably has a polynucleotide sequence which hybridizes under a stringent condition with a polynucleotide consisting of the nucleotide sequence of the nucleotide Nos.289 to 378 in SEQ ID No:2.

**[0058]** Each of the protein (9) and (10), for achieving the function of the protein encoded thereby as a molecule involved in an intracellular signal transduction by a GPCR stimulation, for example, as a G protein  $\alpha$  subunit, usually has a nucleotide sequence of the nucleotide Nos.223 to 399, 859 to 876, 1057 to 1077, 1282 to 1305 in the nucleotide sequence represented by SEQ ID No:2, or preferably has an nucleotide sequence having a homology usually of 85% or more, especially 90% or more with these regions. The total nucleotide number is usually about 963 to 1443, especially 1053 to 1383.

**[0059]** An index indicating which and how many bases in a polynucleotide (10) can be substituted, deleted or added without losing any biological functions of the protein encoded by the polynucleotide (9) is similar to that described above with regard to the first polynucleotide.

<Third polynucleotide>

**[0060]**

(11) A polynucleotide encoding a protein involved in a G protein-coupled receptor mediated signal transduction which comprises at its 5' terminal the nucleotide sequence of the nucleotide Nos.1 to 378 of SEQ ID No:2.

(12) A polynucleotide encoding a protein involved in a G protein-coupled receptor mediated signal transduction which comprises at its 5' terminal a nucleotide sequence having a homology of 70% or more with the polynucleotide consisting of the nucleotide sequence of the nucleotide Nos.1 to 378 of SEQ ID No:2.

**[0061]** In a polynucleotide (12), the nucleotide sequence part corresponding to the 378 nucleotide sequence at the 5' terminal of the protein (11) (hereinafter referred to as a "specific 5' terminal nucleotide sequence") preferably has a homology of 75% or more, especially 80% or more with the specific 5' terminal amino acid sequence of (11).

**[0062]** A polynucleotide (12) preferably has at its 5' terminal a polynucleotide sequence which hybridizes under a stringent condition with the polynucleotide consisting of the nucleotide sequence of the nucleotide Nos.1 to 378 in SEQ ID No:2.

**[0063]** Each of the protein (11) and (12), for achieving the function of the protein encoded thereby as a molecule involved in an intracellular signal transduction by a GPCR stimulation, for example, as a G protein  $\alpha$  subunit, usually has a nucleotide sequence of the nucleotide Nos.223 to 399, 859 to 876, 1057 to 1077, 1282 to 1305 in the nucleotide

sequence represented by SEQ ID No:2, or preferably has a nucleotide sequence having a homology usually of 85% or more, especially 90% or more with these regions. The total nucleotide number is usually about 963 to 1443, especially 1053 to 1383.

**[0064]** An index indicating which and how many bases in a polynucleotide (12) can be substituted, deleted or added without losing any biological functions of the protein encoded by the polynucleotide (11) is similar to that described above with regard to the first polynucleotide.

**[0065]** There is no polynucleotide encoding a G protein  $\alpha$  subunit having the nucleotide sequence of the nucleotide Nos.1 to 378 in SEQ ID No:2 or a sequence analogous thereto.

#### <Inventive polynucleotide production method>

**[0066]** Polynucleotides (7) to (12) can be obtained for example by a screening a human DNA library by a hybridization using as a probe an oligonucleotide (for example, an oligonucleotide of the invention described below) synthesized based on the nucleotide sequence represented by SEQ ID No:2. They can be obtained also by a PCR in a standard manner after preparing suitable primers (for example, oligonucleotides of the invention described below) based for example on the nucleotide sequence of SEQ ID No:2 using as a PCR template a cDNA library for example of a human, rat and mouse. They can be obtained also by a chemical synthesis.

**[0067]** As a cDNA library, one derived from a brain, thymus, testes, spleen, small intestine, uterus and heart is preferred.

**[0068]** A method for obtaining a polynucleotide (10) by introducing a variation into a polynucleotide (9) and a method for obtaining a polynucleotide (12) by introducing a variation into a polynucleotide (11) are as described above.

#### <Application of inventive polynucleotide>

**[0069]** An inventive polynucleotide can be used preferably as a regulator of an intracellular signal transduction mediated by a GPCR stimulation. Typically, it can be used preferably for treating or preventing a disease caused by an abnormality in this intracellular signal transduction.

It is useful especially in treating or preventing a disease associated with an intracellular signal transduction relating to the defect, reduced expression level or reduced function of a protein of the invention.

**[0070]** An inventive polynucleotide can be used preferably also in screening for a substance capable of regulating a signal transduction mediated by a GPCR and a G protein of the invention.

#### <Inventive recombinant vector and transformant>

**[0071]** An inventive recombinant vector is a vector containing an inventive polynucleotide (which herein is a DNA). For example, it may be a vector capable of expressing a protein of the invention.

**[0072]** A vector capable of expressing a protein of the invention can be produced by ligating an inventive polynucleotide to an expressible position downstream of a promoter of an expression vector in accordance with a standard method.

**[0073]** An expression vector may be selected from known vectors capable of replicating in host cells as appropriate depending on the host cells. For example, a pBR322, pUC12, pUC119 and pBluescript can be exemplified when an E.coli is employed as a host cell, while a pUB110 and pC194 are exemplified when a Bacillus organism is employed as a host cell. An Yip5 and Yep24 are exemplified when using a yeast as a host cell. An AcNPV is exemplified when using an insect cell as a host cell. A pUC18 and pUC19 are exemplified when using an animal cell as a host cell.

**[0074]** A host cell may be any of those known in the art without limitation. Those which may be exemplified are bacteria such as an E.coli (for example, K12) and a Bacillus microorganism (for example, MI114), yeasts (for example, AH22), insect cells (for example, Sf cell), animal cells (for example, COS-7 cell, Vero cell, CHO cell and the like).

**[0075]** A method for transforming a host cell with an inventive recombinant vector may be a known method selected suitably depending on the host cell. A known introduction method may for example be a calcium phosphate method, electroporation, lipofection, DEAE dextran method and the like. From transformants, an inventive transformant is selected for example by means of a drug resistance marker possessed by the vector as an index.

#### <Inventive protein production method>

**[0076]** A method for producing a protein of the invention is a method in which an inventive transformant is cultured and an inventive protein is recovered from the resultant culture product.

**[0077]** The conditions of culturing a transformant may be selected appropriately depending on the type of the transformant.

**[0078]** When an inventive transformant is a microorganism, the culture is conducted in a liquid medium or plate medium employed usually for culturing a microorganism. The culture temperature may be within the range allowing a microorganism to be grown, for example from 15 to 40°C. The culture medium pH may also be within the range allowing a microorganism to be grown, for example about pH6 to 8. The culture time period may vary depending on other culturing conditions, and may usually be 1 to 5 days, especially 1 to 2 days. When using an inducible expression vector such as those of the temperature shift type or IPTG inducible type, the induction time period may be within a day, especially within several hours.

**[0079]** Also when an inventive transformant is a mammalian cell, it can be cultured under the condition suitable for said cell. For example, an FBS-supplemented DMEM medium (NISSUI) may be employed to conduct a culture in the presence of 5% CO<sub>2</sub> at a temperature of 36 to 38°C with replacing the medium with a fresh one at an interval of several days. Upon confluent growth, the cells were combined with a trypsin PBS solution to disperse into individual cells and the resultant cell suspension was diluted by several times and inoculated onto a new petri dish, which is then subjected to a subculture. The culture time period is usually 2 to 5 days, especially 2 to 3 days.

**[0080]** Also when an insect cell is employed as a transformant, the culture condition may be adjusted appropriately depending on the type of the cell. For example, an insect cell culture medium such as Grace's medium containing FBS and Yeastlate may be employed to conduct the culture at 25 to 35°C. The culture time period is 1 to 5 days, especially 2 to 3 days. When using as a vector an virus-containing transformant such as a Baculovirus, the culture is continued preferably until the time before the cell death as a result of the onset of the cytoplasmic effect (for example, 3 to 7 days, especially 4 to 6 days).

**[0081]** After completion of the culture, the transformant cells were recovered by a centrifugation, suspended in a suitable buffer if desired, and dispersed by means of a polytron, ultrasonic treatment, homogenizer and the like. The resultant dispersion is centrifuged at about 100 to 150G for about 5 to 10 minutes, and the resultant supernatant is centrifuged at about 18,000 to 20,000G for about 20 to 30 seconds to recover the pellet, whereby obtaining a cell membrane fraction.

**[0082]** The cell membrane fraction thus obtained is subjected to a known protein purification method, such as any of chromatographic methods including ion exchange, hydrophobic, gel filtration and affinity chromatographies, whereby purifying the protein according to the invention.

**[0083]** An inventive protein can be expressed for example as being attached with a histidine tag, or as a glutathion S transferase fusion protein. The former case employs a metal chelate affinity column, while the latter case employs a glutathion S transferase monoclonal antibody column, whereby accomplishing the purification of an inventive protein in a further convenient manner.

<Inventive antisense polynucleotide>

<Aspect>

**[0084]** An inventive antisense polynucleotide is a polynucleotide (13) or (14) shown below.

(13) A polynucleotide which inhibits expression of a protein of the invention which consists of a nucleotide sequence complementary to at least 15 contiguous nucleotides in the nucleotide sequence represented by SEQ ID No:2.

(14) A polynucleotide which inhibits expression of a protein of the invention which hybridizes under an intracellular condition with a polynucleotide consisting of at least 15 contiguous nucleotides in the nucleotide sequence represented by SEQ ID No:2.

**[0085]** An antisense oligonucleotide of the invention hybridizes with a mRNA encoding an inventive protein to inhibit the translation from the mRNA to the protein or cleaves the mRNA, whereby inhibiting the expression of this protein.

**[0086]** While the upper limit of the number of the nucleotides in an antisense polynucleotide is not limited particularly, it is usually about 60 nucleotides for the purpose of achieving the objective.

**[0087]** A polynucleotide (13) preferably has a nucleotide sequence complementary to at least 30 nucleotides, especially at least 50 nucleotides in the polynucleotide represented by SEQ ID No:2. An antisense (14) is preferably one which hybridizes under an intracellular condition also with a polynucleotide consisting of at least 30 nucleotides, especially at least 50 nucleotides in the polynucleotide represented by SEQ ID No:2.

**[0088]** A polynucleotide which hybridizes under an intracellular condition in the invention may for example be a polynucleotide which hybridizes under a stringent condition described below. The stringent condition may for example be a condition involving 2 x SSC, 1 x Denhart's solution at about 60°C.

**[0089]** A polynucleotide (13) preferably has a nucleotide sequence complementary to at least 15 contiguous nucleotides across the both of the region of the nucleotide Nos.1 to 378 and the region of the nucleotide Nos.379 to 1377 in SEQ ID No:2.

**[0090]** Similarly, a polynucleotide (14) is preferably one which hybridizes under an intracellular condition with a polynucleotide consisting of at least 15 contiguous nucleotides across the both of the region of the nucleotide Nos.1 to 378 and the region of the nucleotide Nos.379 to 1377 in SEQ ID No:2.

**[0091]** "At least 15 contiguous nucleotides in the nucleotide sequence represented by SEQ ID No:2" in polynucleotides (13) and (14) which is closer to the 5' terminal of SEQ ID NO:2 is more preferable.

**[0092]** An antisense polynucleotide of the invention may be a single-stranded DNA, double-stranded DNA, single-stranded RNA, double-stranded RNA or DNA-RNA hybrid. When a double-stranded RNA is employed, it is generally called an RNAi. A derivative of such a nucleotide may also be employed as long as it inhibits the expression of an inventive protein. A derivative may for example be a phosphorothioate DNA, H-phosphonate DNA and the like.

**[0093]** Inhibition of an inventive protein expression by an inventive antisense polynucleotide can be verified for example by a method described below. A human brain-derived cell is combined with an antisense at about 5nM to 10  $\mu$ M if necessary together with a known intracellular introduction reagent such as a lipofection reagent, lipofectamine reagent, liposome and the like. Then, from this cell a cell extract is prepared by a known method and an inventive antibody is used to measure expression level of the inventive protein by a known method such as an ELISA or western blotting. This expression level is compared with the level observed in the absence of the antisense.

**[0094]** An inventive antisense polynucleotide may for example be one which reduces the inventive protein expression level for example to 70% or less, preferably to 50% or less based on the level in the absence of the antisense polynucleotide.

**[0095]** An inventive antisense polynucleotide can be produced by a known chemical synthesis method.

<Application of antisense polynucleotide>

**[0096]** An inventive antisense polynucleotide can be used preferably as a regulator of an intracellular signal transduction mediated by a GPCR stimulation. Typically, it can be used preferably for treating or preventing a disease caused by an abnormality in this intracellular signal transduction. It is useful preferably for the purpose especially of suppressing the increased abnormality in this intracellular signal transduction.

<Ribozyme>

<Aspect>

**[0097]** An inventive ribozyme is a ribozyme (15) or (16) shown below.

(15) A ribozyme having an ability of cleaving a polynucleotide of the invention which comprises two polynucleotide regions complementary to two regions respectively consisting of at least 9 contiguous nucleotides which are two regions in the nucleotide sequence represented by SEQ ID No:2.

(16) A ribozyme having an ability of cleaving a polynucleotide of the invention which comprises two polynucleotide regions which hybridizes under an intracellular condition with two regions respectively consisting of at least 9 contiguous nucleotides which are two regions in the nucleotide sequence represented by SEQ ID No:2.

**[0098]** A ribozyme (15) preferably comprises two polynucleotide regions complementary to two regions respectively consisting of at least 10, especially 11 contiguous nucleotides which are two regions in the nucleotide sequence represented by SEQ ID No:2.

**[0099]** A ribozyme (16) preferably comprises two polynucleotide regions which hybridizes under an intracellular condition with two regions respectively consisting of at least 10, especially 11 contiguous nucleotides which are two regions in the nucleotide sequence represented by SEQ ID No:2.

**[0100]** In the ribozymes (15) and (16), the two regions in the nucleotide sequence represented by SEQ ID No:2 may be adjacent to each other, or but may preferably be interrupted by about 1 to 4 nucleotides present between them. For example, a hammer-head ribozyme may contain a single interrupting nucleotide, while a hairpin ribozyme may contain 4 interrupting nucleotides.

**[0101]** A ribozyme is an RNA molecule containing an antisense sequence recognizing a specific site of an RNA, and has an RNA cleavage enzyme activity. As a result, the ribozyme recognizes its target RNA, and cleaves a certain site of the RNA specifically.

**[0102]** An inventive ribozyme may be of a hammer-head or hairpin type. A hammer-head type usually recognize an NUX (N is G, U, C or A, while X is C, U or A) and cleaves a mRNA at the 3'-position of the X.

**[0103]** A hammer-head ribozyme according to the invention may for example be a ribozyme comprising a nucleotide sequence listed below.

A ribozyme comprising the nucleotide sequence:



TCGCCTCCTTCTGATGAGGCCGAAAGGCCGAAACCGCCTCGCGC (SEQ ID No:3).

- 5 The ribozyme having this nucleotide sequence once forms its conformation and then recognizes the nucleotide sequence of the nucleotide Nos.273 to 295 in SEQ ID No:2.  
A ribozyme comprising the nucleotide sequence:

10 CGGCCGCCCCGCTGATGAGGCCGAAAGGCCGAAACTGGGGCCAGC (SEQ ID No:4).

- The ribozyme having this nucleotide sequence once forms its conformation and then recognizes the nucleotide sequence of the nucleotide Nos.111 to 133 in SEQ ID No:2.  
15 A ribozyme comprising the nucleotide sequence:

CAGCGGCCGCCTGATGAGGCCGAAAGGCCGAAACTGTAGCACA (SEQ ID No:5).

- 20 The ribozyme having this nucleotide sequence once forms its conformation and then recognizes the nucleotide sequence of the nucleotide Nos.8 to 30 in SEQ ID No:2.

[0104] A hairpin type ribozyme usually recognizes an NNNG/CN\*GUCNNNNNNNN (N is G, U, C or A), and cleaves a mRNA between N\*G.

- 25 [0105] A hairpin ribozyme according to the invention may for example be a ribozyme comprising a nucleotide sequence listed below.  
A ribozyme comprising the nucleotide sequence:

30 TCGCCTCCTTAGAAGCCTACCAGAGAAACACACGTTGTGGTATATTACCTGGTA

(SEQ ID No:6). The ribozyme having this nucleotide sequence once forms its conformation and then recognizes the nucleotide sequence of the nucleotide Nos.287 to 295 in SEQ ID No:2.

- 35 A ribozyme comprising the nucleotide sequence:

CGGCCGCCCCGAGAAGGGGACCAGAGAAACACACGTTGTGGTATATTACCTGGTA

- 40 (SEQ ID No:7). The ribozyme having this nucleotide sequence once forms its conformation and then recognizes the nucleotide sequence of the nucleotide Nos.116 to 133 in SEQ ID No:2.

A ribozyme comprising the nucleotide sequence:

45 CAGCGGCCGCAAGAAGTAGACCAGAGAAACACACGTTGTGGTATATTACCTGGTA

- (SEQ ID No:8). The ribozyme having this nucleotide sequence once forms its conformation and then recognizes the nucleotide sequence of the nucleotide Nos.12 to 30 in SEQ ID No:2.

[0106] While an inventive ribozyme usually consists of an RNA, a one which includes a deoxyribonucleotide or a derivative such as a phosphorthioate DNA which is difficult to be decomposed in vivo are also included in the inventive ribozyme.

- 55 [0107] An inventive ribozyme can be produced by a known chemical synthesis method, in vitro or in vivo transcription and the like. Typically, an in vitro transcription involves the ligation of a DNA having a sequence complementary to the sequence of SEQ ID No:3, 4, 5, 6, 7 or 8 to the downstream of the DNA encoding a promoter such as T7, T3 or SP6. Using this DNA as a template, a transcription reaction by an RNA polymerase is conducted. The resultant transcription product can be used as an RNA for the ribozyme.

**[0108]** An in vivo transcription involves the integration of a DNA having a sequence complementary to the sequence of SEQ ID No:3, 4, 5, 6, 7 or 8 into an mammalian expression vector followed by the transduction of this expression vector into a mammalian cell. As a result of the cellular transcription mechanism, an RNA having the sequence of SEQ ID No:3, 4, 5, 6, 7 or 8 is synthesized.

<Application of ribozyme>

**[0109]** An inventive ribozyme can be used preferably as a regulator of an intracellular signal transduction mediated by a GPCR stimulation. Typically, it can be used preferably for treating or preventing a disease caused by an abnormality in this intracellular signal transduction. It is useful preferably for the purpose especially of suppressing the increased abnormality in this intracellular signal transduction.

<Inventive antibody>

<Aspect>

**[0110]** An inventive antibody is an antibody which recognizes an inventive protein specifically. The inventive antibody may be a polyclonal antibody or monoclonal antibody. The inventive antibody includes an antibody having an antigen binding ability toward a polypeptide consisting of 5 contiguous amino acids, preferably 10 amino acids in the amino acid sequence constituting an inventive protein. In addition, a derivative of such an antibody (chimera antibody and the like) or a one labeled with an enzyme such as a peroxidase are also included in the inventive antibody.

<Method for producing inventive antibody>

**[0111]** Any of the antibodies described above can be produced in accordance with a known production method (for example, Current protocols in Molecular Biology edit. Ausubel et al. (1987) Publish. John Wiley and Sons. Section 11.12-11.13).

**[0112]** Typically, when an inventive antibody is a polyclonal antibody, it can be obtained by immunizing a non-human animal such as a rodent animal with an inventive protein followed by the isolation from the serum of this immunized animal in accordance with a standard method. When an inventive antibody is a monoclonal antibody, it can be obtained from a hybridoma produced by immunizing a non-human animal such as a mouse with a polypeptide having an inventive protein or its partial sequence followed by fusing the spleen cell of this immunized animal with a myeloma cell (Current protocols in Molecular Biology edit. Ausubel et al. (1987) Publish. John Wiley and Sons. Section 11.4-11.11).

<Application of antibody>

**[0113]** An inventive antibody can be used preferably as a regulator of an intracellular signal transduction mediated by a GPCR stimulation. Typically, it can be used preferably for treating or preventing a disease caused by an abnormality in this intracellular signal transduction. It is useful preferably for the purpose especially of suppressing the increased abnormality in this intracellular signal transduction.

**[0114]** An inventive antibody can preferably be used also in the affinity chromatography for purifying an inventive protein as well as in the screening for a substance which may affect the expression of an inventive protein.

<Inventive oligonucleotide>

**[0115]** An inventive oligonucleotide is an oligonucleotide (17) or (18) shown below.

(17) An oligonucleotide capable of recognizing a polynucleotide represented by SEQ ID NO:2 specifically which consists of at least 17 contiguous nucleotides in the nucleotide sequence represented by SEQ ID No:2.

(18) An oligonucleotide capable of recognizing a polynucleotide represented by SEQ ID NO:2 specifically which has a homology of 80% or more with at least 17 contiguous nucleotides in the nucleotide sequence represented by SEQ ID No:2.

**[0116]** The length of each of the oligonucleotides (17) and (18) can be selected appropriately depending on the use. It may be the full length of SEQ ID No:2.

**[0117]** An oligonucleotide (18) preferably has a homology of 85% or more, especially 90% or more with an oligonucleotide (17).

**[0118]** The expression "recognize specifically" in conjunction with oligonucleotides (17) and (18) means that each

oligonucleotide can be employed to detect the nucleotide sequence represented by SEQ ID No:2 specifically or selectively by a known nucleotide sequencing means such as a northern blotting or PCR.

[0119] An inventive oligonucleotide can be used as a probe or primer which can detect or amplify an RNA generated as a result of the expression of an inventive DNA or a polynucleotide derived therefrom in a specific manner. Typically, it can be used as a probe or primer in a known method for detecting a certain nucleotide sequence, such as a Northern blotting, in situ hybridization or PCR.

[0120] As a result, the absence or presence of the expression of, or the expression level of an inventive polynucleotide can be assessed. Accordingly, an inventive oligonucleotide can preferably be used for diagnosing a disease caused by a signal transduction abnormality resulting from the defect of and the abnormal increase or decrease in the expression level of the inventive protein. A test sample may be a total RNA prepared by a standard method from a sample taken from a tissue of a subject such as an uterus or any of various polynucleotides prepared from such an RNA.

[0121] An inventive oligonucleotide may for example be ones having the nucleotide sequences of

5' -ATGGGTCTGTGCTACAGTCTGCGG (SEQ ID No:9)

and

5' -ACGATGGTGCTTTTCCCAGACTCACCAGCCCCGAGCA (SEQ ID No:10).

This oligonucleotide set can preferably be used as PCR primers for amplifying a protein of the invention.

<Inventive screening method>

[0122] A polynucleotide of the invention can be used for screening for a substance which activates or inhibits (or suppress) the cellular signal transduction mediated by a GPCR and an inventive protein.

[0123] A GPCR stimulating signal is transmitted to an effector via the activation of a G protein as a result of the GDP/GTP exchange reaction on a G protein  $\alpha$  subunit. Accordingly, by using the change in the activity of this effector as an index in screening test substances, a substance capable of activating or inhibiting the signal transduction mediated by a GPCR and a G protein  $\alpha$  subunit can be identified. In addition, also by using the change in the level of the binding of a GTP to a membrane fraction of a cell expressing a G protein as an index in screening test substances, a substance capable of activating or inhibiting the signal transduction mediated by a GPCR and a G protein  $\alpha$  subunit can be identified.

[0124] Otherwise, by screening test substances which alter the level of the expression of an inventive protein, a substance capable of activating or inhibiting the signal transduction mediated by a GPCR and a G protein  $\alpha$  subunit can be identified.

<First method>

[0125] A method for screening for a substance capable of regulating a signal transduction mediated by a GPCR and an inventive protein comprising:

(a) a step for bringing a test substance into contact with a test cell having a recombinant vector containing an inventive polynucleotide (which herein is a DNA encoding an inventive protein) and a recombinant vector containing a DNA encoding a GPCR;

(b) a step for measuring the G protein effector activity or the index value correlating therewith in the test cell (hereinafter abbreviated as "effector activity"); and

(c) a step for comparing this effector activity with the effector activity in the test cell which has not been brought into contact with the test substance, whereby selecting a test substance capable of altering the effector activity in the test cell.

[0126] In the first method, an agonist of GPCR can be selected.

[0127] A cell in which a recombinant vector having a DNA encoding an inventive protein and a recombinant vector having a DNA encoding a GPCR are contained may for example be but not limited to a mammalian cell or insect cell. A mammalian cell may be any known cell such as a Vero cell, Hela cell, CV1 cell, COS1 cell, CHO cell and the like,

which may be employed without limitation. An insect cell may be any known cell such as a Sf cell, MG1 cell, High Five™ cell, and the like, which may be employed without limitation. The type of the vector is not limited particularly, and any known vector may be selected depending on the type of the cell.

**[0128]** A DNA encoding a GPCR can be obtained by a method for screening a human cDNA library using a probe designed based on the nucleotide sequences described in a GPCR database ([www.cmbi.kun.nl/7tm/](http://www.cmbi.kun.nl/7tm/)), by a method for conducting a PCR using as a template a human cDNA library together with the primers designed based on the nucleotide sequences described above, or by a chemical synthesis method and the like.

**[0129]** An effector of a G protein may be an effector which is a target of a G protein  $\alpha$  subunit or may be an effector which is a target of a G protein  $\beta\gamma$  subunit. It is also possible to measure an index value correlating with the effector activity of a G protein.

**[0130]** An effector which is targeted by a G protein  $\alpha$  subunit may for example be an adenylyl cyclase,  $\text{Ca}^{2+}$  channel,  $\text{K}^{+}$  channel, phospholipase  $\text{C}\beta$  and the like. The adenylyl cyclase activity be assessed by measuring the intracellular cAMP level. The  $\text{Ca}^{2+}$  channel activity can be assessed by measuring the cell membrane electric potential. The  $\text{K}^{+}$  channel activity can be assessed by measuring the cell membrane electric potential. The phospholipase  $\text{C}\beta$  activity can be assessed by measuring the  $\text{Ca}^{2+}$  level.

**[0131]** An effector which is targeted by a G protein  $\beta\gamma$  subunit may for example be an adenylyl cyclase,  $\text{Ca}^{2+}$  channel,  $\text{K}^{+}$  channel, phospholipase  $\text{C}\beta$ , phosphatidylinositol 3-kinase  $\beta$  or  $\gamma$  and the like. The phosphatidylinositol 3-kinase  $\beta$  or  $\gamma$  activity can be assessed by measuring the  $\text{Ca}^{2+}$  level.

**[0132]** An effector to be examined for its activity is preferably an effector targeted by a G protein  $\alpha$  subunit, with an adenylyl cyclase being more preferred. The intracellular cAMP level which reflects the adenylyl cyclase activity can be measured by a known method such as an RIA employing an anti-cAMP antibody obtained by immunizing a mouse, rat, rabbit, goat, cattle and the like together with a  $^{125}\text{I}$ -labeled cAMP, other EIA employing a combination of an anti-cAMP antibody and a labeled cAMP, a SPA method employing a scintillant obtained by immobilizing an anti-cAMP antibody using a protein A or an antibody against an animal IgG used for producing an anti-cAMP antibody together with a  $^{125}\text{I}$ -labeled cAMP, an EFC method employing a combination of an anti-cAMP antibody, enzyme donor-binding cAMP and enzyme blank acceptor, and the like. Any of these measurements can be accomplished using a commercial kit.

**[0133]** An intracellular cAMP level can be assessed also by a method in which, for example, a CRE (cAMP response element, which reacts with a cAMP)-containing DNA is inserted into the upstream of the reporter gene of a reporter gene vector to form a CRE-reporter gene vector, and this vector is also introduced into a test cell and then the reporter gene expression level is measured. In a cell into which a CRE-reporter gene vector has been introduced, a stimulation accompanied with an elevation in the cAMP level induces a reporter gene expression mediated by the CRE and the subsequent reporter protein production. On the contrary, a reduction in the cAMP level leads to a reduction in the CRE-mediated reporter protein production. By using a CRE reporter gene vector, the cAMP level can be measured conveniently at a high sensitivity.

**[0134]** As a reporter gene, any of known genes can be employed without limitation, including a luciferase gene, secretor alkaline phosphatase (SEAP) gene, chloramphenicol acetyltransferase (CAT) gene,  $\beta$ -galactosidase gene and the like. Any of these genes can be examined for its expression level using a commercial measurement kit described below. The luciferase gene expression level can be measured by adding a luminescent substrate luciferin (manufactured for example by TOYO INK) to a cell solution followed by measuring the luminescence resulting from the decomposition of the substrate using a luminometer, liquid scintillation counter or top counter. Expression level of the alkaline phosphatase gene can be determined for example by using  $\text{L}\mu\text{Mi-Phos530}$  (WAKO PURE CHEMICAL). Expression level of the chloramphenicol acetyltransferase gene can be determined using a FAST CAT Chloramphenicol Acetyltransferase Assay Kit (WAKO PURE CHEMICAL). Expression level of the  $\beta$ -galactosidase gene can be determined using an AURORA Gal-XE (WAKO PURE CHEMICAL).

**[0135]** In the case for example where a luciferase gene is employed, a CRE-containing DNA is inserted into a multiple cloning site in the upstream of a luciferase gene such as a PICK-A-GENE Basic Vector or a PICK-A-GENE Enhancer vector (TOYO INK) and the like, which is then used as a CRE reporter gene vector.

**[0136]** The type of a test substance is not limited particularly. Those which may be exemplified are proteins, peptides, non-peptide compounds (nucleotides, amines, saccharides, lipids and the like), organic low molecular compounds, inorganic low molecular compounds, fermentation products, cell extracts, plant extracts, animal tissue extracts and the like.

**[0137]** The contact of a cell with a test substance may be effected under a condition avoiding the cell death and allowing an inventive protein and a GPCR to be expressed from an introduced vector (temperature, pH, medium composition). The concentration of a test substance upon contact with a cell may for example be about 0.001 to 10  $\mu\text{M}$ , although it may vary depending on the type of the substance.

**[0138]** A test substance which increase the effector activity of a test cell brought into contact with the test substance for example by about 25%, preferably about 50%, more preferably about 100%, when compared with the effector

activity in the test cell which was not brought into contact with the test substance, can be selected as a GPCR agonist.

<Second method>

5 **[0139]** A method for screening for a substance capable of regulating a signal transduction mediated by a GPCR and an inventive protein comprising:

- 10 (a) a step for bringing a test substance into contact with a test cell having a recombinant vector containing an inventive polynucleotide (which herein is a DNA encoding an inventive protein) and a recombinant vector containing a DNA encoding a GPCR;
- (b) a step for measuring the G protein effector activity in the test cell; and
- (c) a step for comparing this effector activity with the effector activity when the said test substance has been brought into contact with a control cell having no recombinant vector containing a DNA encoding an inventive protein but having a recombinant vector containing a DNA encoding a GPCR, whereby selecting a test substance causing a difference in the effector activity between the test cell and the control cell.

**[0140]** In the second method, a test substance which gives the effector activity of a test cell having an inventive protein expression vector which is higher than the effector activity in a control cell having no such a vector may be searched for. As a result, a substance which activates any stage of the signal transduction mediated by a GPCR and the inventive protein can be selected as a candidate compound.

**[0141]** A test substance which increase the effector activity of a test cell for example by about 20%, preferably about 50%, more preferably about 100%, when compared with a control cell can be selected as a signal transduction activator.

**[0142]** Otherwise, the aspect is similar to that in of the first method discussed above.

25 <Third method>

**[0143]** A method for screening for a substance capable of regulating a signal transduction mediated by a GPCR and an inventive protein comprising:

- 30 (a) a step for bringing a test substance into contact with a test cell having a recombinant vector containing an inventive polynucleotide (which herein is a DNA encoding an inventive protein) and a recombinant vector containing a DNA encoding a GPCR;
- (b) a step for measuring the G protein effector activity in the test cell; and
- 35 (c) a step for comparing this effector activity with the effector activity when the said test substance has been brought into contact with a control cell having no recombinant vector containing a DNA encoding a GPCR but having a recombinant vector having a DNA encoding an inventive protein, whereby selecting a test substance causing a difference in the effector activity between the test cell and the control cell.

**[0144]** In the third method, a test substance which gives the effector activity of a test cell having a GPCR expression vector which is higher than the effector activity in a control cell having no GPCR expression vector may be searched for. As a result, a GPCR agonist can be selected as a candidate substance.

**[0145]** A test substance which increase the effector activity of a test cell for example by about 20%, preferably about 50%, more preferably about 100%, when compared with a control cell can be selected as a signal transduction activator.

**[0146]** Otherwise, the aspect is similar to that in of the first method discussed above.

45 <Fourth method>

**[0147]** A method for screening for a substance capable of regulating a signal transduction mediated by a GPCR and an inventive protein comprising:

- 50 (a) a step for bringing a test substance and a GPCR ligand into contact with a test cell having a recombinant vector containing an inventive polynucleotide (which herein is a DNA encoding an inventive protein) and a recombinant vector containing a DNA encoding a GPCR protein;
- (b) a step for measuring the G protein effector activity in the test cell; and
- 55 (c) a step for comparing this effector activity with the effector activity in the test cell which has not been brought into contact with the test substance but has been brought into contact with the ligand, whereby selecting a test substance capable of altering the effector activity in the test cell.

**[0148]** In the fourth method, a test substance which gives increased or reduced effector activity in the control cell which has not been brought into contact with the test substances when compared with the effector activity in the test cell which has been brought into contact with the test substances may be searched for. As a result, a substance which activates or inhibits any stage of the signal transduction initiated from the binding of a GPCR ligand to a GPCR can be selected, including a GPCR agonist or antagonist.

**[0149]** A GPCR ligand may for example be amine molecules. It is preferable especially to use dopamine. The ratio between a ligand to be brought into contact with a cell and a test substance, when represented as the molar ratio of ligand:test substance, may for example be about 1:0.1 to 1:100, preferably about 1:1 to 1:50.

**[0150]** The percentage effector activity in a test cell which has been brought into contact with the both of a GPCR ligand and a test substance is calculated, on the bases of the effector activity in the test cell which has been brought into contact only with the GPCR ligand being regarded as 100% and the effector activity in the test cell which has been brought into contact with none of the GPCR ligand or the test substance as 0%. A test substance which gives a % effector activity in a test cell which has been brought into contact with the both of a GPCR ligand and the test substance of 85% or less, preferably 70% or less, especially 50% or less can be selected as a candidate of the cellular signal transduction inhibitor or suppressor. On the other hand, a test substance which raises this percentage to 125% or more, preferably 150% or more, especially 200% or more can be selected as a candidate of the cellular signal transduction activator.

**[0151]** Otherwise, the aspect is similar to that in of the first method discussed above.

<Fifth method>

**[0152]** A method for screening for a substance capable of regulating a signal transduction mediated by a GPCR and an inventive protein comprising:

(a) a step for bringing a test substance and a GPCR ligand into contact with a test cell having a recombinant vector containing an inventive polynucleotide (which herein is a DNA encoding an inventive protein) and a recombinant vector containing a DNA encoding a GPCR protein;

(b) a step for measuring the G protein effector activity in the test cell;

(c) a step for comparing this effector activity with the effector activity in the test cell which has not been brought into contact with the test substance but has been brought into contact with the ligand, whereby investigating the change in the effector activity in the test cell; and

(d) a step for comparing the rate of change in this effector activity with the rate of change in the effector activity when the said test substance and said ligand has been brought into contact with a control cell having no recombinant vector containing a DNA encoding a GPCR but having a recombinant vector containing a DNA encoding an inventive protein, whereby selecting a test substance causing a difference in the rate of change in the effector activity between the test cell and the control cell.

**[0153]** In the fifth method, a test substance which gives an elevated rate of change in the effector activity in a test cell having a recombinant vector containing a DNA encoding a GPCR protein when compared with the rate of change in the effector activity in a control cell having no such a vector may be searched for. As a result, a substance which serves as an antagonist against an exogenous GPCR can be selected. While in the fourth method described above an antagonist against an endogenous GPCR is also selected, by subjecting the substances obtained by the fourth method to a screening by the fifth method, a substance serving as an antagonist against the endogenous GPCR can selectively be eliminated.

**[0154]** A test substance which increase the rate of change in the effector activity of a test cell for example by about 15%, preferably about 30%, more preferably about 50%, when compared with a control cell can be selected as a candidate of an exogenous GPCR-directed antagonist.

**[0155]** Otherwise, the aspect is similar to that in of the first method discussed above.

<Sixth method>

**[0156]** A method for screening for a substance capable of regulating a signal transduction mediated by a GPCR and an inventive protein comprising:

(a) a step for bringing a test substance into contact with a cell membrane fraction of a cell having a recombinant vector containing an inventive polynucleotide (which herein is a DNA encoding an inventive protein) and a cell membrane fraction of a cell having a recombinant vector containing a DNA encoding a GPCR, or a cell membrane fraction of a cell having the recombinant vector containing a polynucleotide encoding an inventive protein (which

herein is a DNA encoding an inventive protein) and the recombinant vector containing the DNA encoding the GPCR;  
 (b) a step for assaying the level of the binding of GTP to the cell membrane fraction; and  
 (c) a step for comparing the assayed level of this GTP binding with the assayed level of the GTP binding to the  
 cell membrane fraction which has not been brought into contact with the test substance, whereby selecting a test  
 substance capable of altering the assayed level of the GTP binding to the cell membrane fraction.

**[0157]** When a cell expressing a GPCR and a G protein is stimulated by a GPCR ligand, then a GTP is bound to a G protein  $\alpha$  subunit. This phenomenon is observed also in a membrane fraction of a cell which expresses a GPCR and a G protein. Accordingly, in the sixth method, a substance which increases the level of the binding of a GTP to this cell membrane fraction can be selected as a GPCR agonist.

**[0158]** Usually, a GTP bound to a G protein  $\alpha$  subunit is decomposed into a GDP. Accordingly, a GTP analogue which is capable of binding to an inventive protein but is not decomposed by a GTPase is used to measure the level of the binding of this GTP analogue to an inventive protein, whereby assaying the level of the binding of the GTP to the inventive protein. Such a GTP analogue may for example be a GTP $\gamma$ S, G $_{pp}$ NH $_p$  and the like.

**[0159]** For measuring the level of the binding of a GTP analogue to a cell membrane fraction, the GTP analogue is labeled for example with a radiolabel, and then the labeled GTP analogue is added to the cell membrane fraction and incubated for a certain period, and then the radioactivity in the cell membrane fraction is measured by a scintillation counter and the like.

**[0160]** The methods for preparing and characterizing a cell membrane fraction are as described above.

**[0161]** A test substance which increases a level of the binding of a GTP to a cell membrane fraction in a test cell which has been brought into contact with a test substance for example by about 25%, preferably about 50%, more preferably about 100%, when compared with a test cell which has not been brought into contact with a test substance can be selected as a candidate GPCR agonist.

**[0162]** Otherwise, the aspect is similar to that in of the first method discussed above.

<Seventh method>

**[0163]** A method for screening for a substance capable of regulating a signal transduction mediated by a GPCR and an inventive protein comprising:

(a) a step for bringing a test substance and a GPCR ligand into contact with a cell membrane fraction of a cell having a recombinant vector containing an inventive polynucleotide (which herein is a DNA encoding an inventive protein) and a cell membrane fraction of a cell having a recombinant vector containing a DNA encoding a GPCR, or a cell membrane fraction of a cell having the recombinant vector containing a polynucleotide encoding an inventive protein (which herein is a DNA encoding an inventive protein) and the recombinant vector containing the DNA encoding the GPCR;

(b) a step for assaying the level of the binding of GTP to this cell membrane fraction; and

(c) a step for comparing the assayed level of this GTP binding with the assayed level of the GTP binding in the cell membrane fraction which has not been brought into contact with the test substance but has been brought into contact with said ligand, whereby selecting a test substance capable of altering the assayed level of the GTP binding to the cell membrane fraction.

**[0164]** In the seventh method, a substance which activates or inhibits (or suppresses) any stage (through the time of the binding of a GTP to a G protein  $\alpha$  subunit) in a cellular signal transduction can be searched for, including GPCR receptor agonist and antagonist.

**[0165]** The percentage radioactivity in a membrane fraction which has been brought into contact with the both of a GPCR ligand and a test substance is calculated, on the bases of the radioactivity in the membrane fraction which has been brought into contact only with the GPCR ligand being regarded as 100% and the radioactivity in the membrane fraction which has been brought into contact with none of the GPCR ligand or the test substance as 0%. A test substance which gives a % radioactivity when brought into contact with the both of a GPCR ligand and the test substance of 75% or less, preferably 50% or less, especially 25% or less can be selected as a candidate of the cellular signal transduction inhibitor or suppressor. On the other hand, a test substance which raises the percentage radioactivity when brought into contact with the both of a GPCR ligand and the test substance to 125% or more, preferably 150% or more, especially 200% or more can be selected as a candidate of the cellular signal transduction activator.

**[0166]** Otherwise, the aspect is similar to that in of the sixth method discussed above. A ligand which can be employed and the ratio between the ligand and a test substance are similar to those in the fourth method.

## &lt;Eighth method&gt;

**[0167]** A method for screening for a substance capable of regulating a signal transduction mediated by a GPCR and an inventive protein comprising:

- (a) a step for bringing a test substance into contact with a test cell capable of expressing a protein of the invention;
- (b) a step for measuring the expression level of the protein of the invention in the test cell; and
- (c) a step for comparing this expression level with the expression level of said protein in the test cell which has not been brought into contact with the test substance, whereby selecting a test substance capable of altering the expression level of said protein in the test cell.

**[0168]** In this method, a substance which activates or inhibits a signal transduction mediated by an inventive protein by increasing or reducing the expression of the inventive protein can be selected.

**[0169]** The expression level of an inventive protein can be determined by measuring the level of the corresponding mRNA. The level of this mRNA can be determined by a known method such as a Northern blotting using an inventive probe described above or a PCR using inventive primers described above.

**[0170]** Specifically, a Northern blotting can be conducted in such a manner that an RNA is prepared from a test cell by a standard method, transferred onto a nylon membrane and the like, hybridized with a probe labeled for example with a radioisotope or fluorescent substance, and then a double strand of the probe with the RNA is detected by a method suitable for the label. A PCR can be conducted in such a manner that a cDNA is prepared from a mRNA of a test cell and used as a template to perform a PCR by a standard method using an inventive oligonucleotide set as primers.

**[0171]** The expression level of an inventive protein can be determined by quantifying the protein directly. The level of this protein can be determined by a known method such as a Western blotting using an inventive antibody.

**[0172]** As a cell expressing an inventive protein, a mammalian cell, preferably a human cell is employed. It is preferred particularly to use a cell derived from a human brain, thymus, testes, spleen, small intestine, uterus and heart.

## &lt;Inventive screening kit&gt;

**[0173]** A first inventive kit for screening for a substance capable of regulating a signal transduction mediated by a GPCR and a protein according to the invention comprises a test cell having a recombinant vector containing an inventive polynucleotide (which herein is a DNA) and a reagent for measuring the G protein effector activity.

**[0174]** This kit can be used in the first inventive screening method described above. A test cell and the reagents for measuring a G protein effector activity are as described above. For performing the first screening method using this kit, the test cell is transduced with a recombinant vector having a DNA encoding a GPCR independently. Alternatively, the test cell may be one having a GPCR expression vector in addition to a recombinant vector having an inventive polynucleotide.

**[0175]** Furthermore, the first inventive screening kit may contain a control cell which does not have a recombinant vector having an inventive polynucleotide (which herein is a DNA) but has a control cell having a recombinant vector having a DNA encoding a GPCR. In such a case, it can be used in the second inventive screening method.

**[0176]** Moreover, the first inventive screening kit may contain a control cell which does not have a recombinant vector having a DNA encoding a GPCR but has an inventive polynucleotide (which herein is a DNA). In such a case, it can be used in the third inventive screening method.

**[0177]** The first inventive screening kit may further contain a GPCR ligand. In such a case, it can be used in the fourth inventive screening method. A ligand is employed also as described above.

**[0178]** The first inventive screening kit may further contain a GPCR ligand and a control cell having no recombinant vector having a DNA encoding a GPCR but having a recombinant vector having an inventive polynucleotide (which herein is a DNA). In such a case, it can be used in the fifth inventive screening method.

**[0179]** The second inventive kit for screening for a substance capable of regulating a signal transduction mediated by a GPCR and a protein according to the invention comprises a cell having a recombinant vector containing an inventive polynucleotide (which herein is a DNA); and a GTP analogue which can bind to the protein of the invention but can not be cleaved by a GTPase. This kit can be employed in the sixth inventive screening method. A GTP analogue is employed also as described above.

**[0180]** For performing the sixth screening method using this kit, a cell is transduced with a recombinant vector having a DNA encoding a GPCR independently. Alternatively, such a cell may be one having a GPCR expression vector in addition to a recombinant vector having an inventive DNA.

**[0181]** The third inventive kit for screening for a substance capable of regulating a signal transduction mediated by a GPCR and a protein according to the invention comprises a cell having a recombinant vector containing an inventive



polynucleotide (which herein is a DNA) and a GPCR expression vector; and a GTP analogue which can bind to the protein of the invention but can not be cleaved by a GTPase. This kit can be employed in the sixth inventive screening method.

**[0182]** Each of the second and third screening kits according to the invention may further contain a GPCR ligand. In such a case, it can be used in the 7th inventive screening method.

**[0183]** The fourth inventive kit for screening for a substance capable of regulating a signal transduction mediated by a GPCR and a protein according to the invention comprises a cell capable of expressing a protein of the invention; as well as an inventive probe, inventive primers or inventive antibody. This kit can be used in the 8th inventive screening method.

#### <Pharmaceuticals>

**[0184]** A protein and an antibody according to the invention can be used as pharmaceuticals by being administered in an effective amount to a mammal such as a human in the forms described below.

**[0185]** A protein and an antibody according to the invention can be formulated into pharmaceutical composition in a mixture with inactive carriers, such as pharmaceutically acceptable carriers (including excipient, extender, binder, lubricant and the like) as well as customary additives. Such a pharmaceutical composition may be given orally or parenterally depending on the dosage form (oral formulation such as tablet, pill, capsule, powder, granule, syrup and the like; parenteral formulation such as injection formulation, drip infusion formulation, dermal formulation, suppository and the like). The dose may vary depending on the type of the active ingredient, administration route, subject and the age, body weight and conditions of the patient, and may be about 0.01 to 100 mg a day, which can be given all at once or in several portions.

**[0186]** A polynucleotide, antisense polynucleotide and ribozyme of the invention can be administered in an effective amount to a mammal such as a human as pharmaceuticals in the forms of the pharmaceutical compositions described above. Otherwise, they can be introduced into a cell of a subject utilizing a liposome delivery system employing a liposome in which a drug to be delivered is encapsulated, a microinjection method, a direct injection method, a gene gun and the like. Also in such cases, the dose and the administration mode can be selected appropriately by those skilled in the art, although it may vary depending on the age, body weight and conditions of the patient.

**[0187]** An inventive polynucleotide can be introduced into a target cell also by integrating into a virus vector for a gene therapy.

**[0188]** A substance capable of regulating a signal transduction obtained by an inventive screening method can be administered in an effective amount to a mammal such as a human in the forms of the pharmaceutical compositions described above. When this substance is one encoded by a DNA, it can be introduced into a target cell also by integrating into a virus vector for a gene therapy.

#### EFFECT OF THE INVENTION

**[0189]** According to the invention, a protein which can be regarded as a novel G protein involved in a cellular signal transduction and a polynucleotide encoding the same are provided.

**[0190]** Moreover, an inventive protein can be regarded as a protein involved in a signal transduction mediated by a GPCR and a G protein involved in the differentiation and the proliferation of a cell, since the G protein effector activity in a cell having vectors expressing the GPCR and the inventive protein respectively is higher than the relevant effector activity in a cell having no vector expressing the inventive protein.

**[0191]** Furthermore, an inventive protein is considered to be one of G proteins, since it has regions having a high homology with the amino acid sequence conserved as a GTP binding site and a GTPase activation site among G proteins and the amino acid sequence of a trimer forming domain conserved among G proteins.

**[0192]** Accordingly, an inventive protein and a polynucleotide encoding the same can preferably be employed as a regulator of an intracellular signal transduction mediated by a GPCR and the inventive protein. Moreover, it is useful in treating or preventing a disease caused by the abnormality in this cellular signal transduction. Specifically, it can preferably be employed for treating or preventing a disease caused by an intracellular signal transduction due to the defect, reduced expression level or reduced function of a protein of the invention.

**[0193]** Moreover, an inventive polynucleotide can preferably be used in the screening for a substance capable of regulating a signal transduction mediated by a GPCR stimulation.

**[0194]** Inventive antibody, antisense and ribozyme are employed preferably as regulators of an intracellular signal transduction mediated by a GPCR and an inventive protein. In addition, they can be used preferably in treating or preventing a disease caused by an abnormality in this intracellular signal transduction. They are useful preferably for the purpose especially of suppressing the increased abnormality in this intracellular signal transduction.

**[0195]** Furthermore, an antibody of the invention can preferably be used also in the affinity chromatography for pu-

rifying an inventive protein as well as in the screening for a substance which may affect the expression of an inventive protein.

**[0196]** A substance obtained by a screening method of the invention can be used as a regulator of a signal transduction mediated by a GPCR and an inventive protein. In addition, it can be used preferably in treating or preventing a disease caused by an abnormality in this intracellular signal transduction.

**[0197]** An inventive oligonucleotide can preferably be used in diagnosing of a disease caused by an intracellular signal transduction due to the defect of, and abnormally increased or reduced expression level of a protein of the invention.

**[0198]** Since an inventive protein is found to be expressed in the tissue of a brain such as a cerebrum, the inventive protein, polynucleotide, antibody, antisense, ribozyme and a substance obtained by an inventive screening method are considered to be useful in preventing or treating a neuropathy and the like. An inventive oligonucleotide is also considered to be useful in diagnosing a neuropathy.

**[0199]** Also since an inventive protein is found to be expressed in a heart, the inventive protein, polynucleotide, antibody, antisense, ribozyme and a substance obtained by an inventive screening method are considered to be useful in preventing or treating a cardiac disease and the like (Targets, 2002, vol.1, p206-213).

**[0200]** Also since an inventive protein is found to be expressed in a thymus and a spleen, the inventive protein, polynucleotide, antibody, antisense, ribozyme and a substance obtained by an inventive screening method are considered to be useful in preventing or treating an immune disease and the like.

## EXAMPLES

**[0201]** The present invention is further described in the following Examples, which are not intended to restrict the invention.

**[0202]** In the following Examples, an inventive protein is sometimes abbreviated as "Gm1".

(Example 1) Cloning of cDNA encoding human Gm1 protein

**[0203]** A pCR-Gm1 which is a plasmid comprising a DNA encoding a full-length human Gm1 was prepared as described below.

20 ng of a plasmid DNA from a human brain-derived cDNA library (Takara) (pAP3neo) was employed as a template together with 10  $\mu$ M of a forward primer: prGm1ATG(5'-ATGGGTCTGTGCTACAGTCTGCGG; SEQ ID No: 11) and 10  $\mu$ M of a reverse primer prGNAL3'(5'-TCACAAGAGCTCATACTGCTT; SEQ ID No:12) as well as TAKARA LA Taq polymerase (TAKARA LA Taq with GC Buffer, Takara) to perform a PCR to obtain an amplified DNA.

**[0204]** The PCR condition involved 35 cycles, each cycle involving incubations at 95°C for 30 seconds followed by 60°C for 30 seconds followed by 72°C for 2 minutes.

**[0205]** The resultant DNA was subjected to an agarose gel electrophoresis followed by a purification with a QIAquick Gel Extraction kit (QIAGEN), and then recovered. This purified and recovered DNA was used as an insert DNA.

**[0206]** Subsequently, a TOPO TA Cloning Kit (Invitrogen) was used and the attached protocol was followed to insert the insert DNA(50ng) into a cloning site of a pCR2.1-TOPO vector (10ng), whereby obtaining a pCR-Gm1.

**[0207]** The DNA thus obtained was subjected to an ABI377 DNA sequencer to determine the nucleotide sequence, and was revealed to contain the nucleotide sequence of the nucleotide Nos.1-1377 in the nucleotide sequence represented by SEQ ID NO:2 and to encode the full-length amino acid sequence represented by SEQ ID NO:1.

(Example 2) Detection of expression profile of nucleic acid encoding Gm1

**[0208]** In order to amplify a nucleic acid encoding Gm1 specifically, a forward primer prGmlrt-5' (5'-ATGGGGTGTGTTT-GGGCGGCAACA; SEQ ID No:13) and a reverse primer prGmlrt-3' (5' -ACGATGGTGCTTTCCAGACTCAC-CAGCCCCGAGCA; SEQ ID No:14) were produced. Each 1  $\mu$ g of human bone marrow-derived total RNA (Ambion), human brain-derived total RNA (Ambion), human spleen-derived total RNA (Ambion), human thymus-derived total RNA (Ambion), human small intestine-derived total RNA (Ambion), human liver-derived total RNA (Ambion), human placenta-derived total RNA (Ambion), human cervix-derived total RNA (Ambion), human uterus-derived total RNA (Ambion), human heart-derived total RNA (Ambion), human skeletal muscle-derived total RNA (Ambion), human testis-derived total RNA (Ambion) and human kidney-derived total RNA (Ambion) were employed as templates together with each 10  $\mu$ M of the primer set described above and a Superscript One-Step RT-PCR System (Invitrogen) and the attached protocol was followed to conduct a RT-PCR to amplify a mRNA. The condition of the RT-PCR involved an incubation at 55°C for 30 minutes for a reverse transcription reaction, followed by 35 cycles, each cycle involving incubations at 94°C for 20 seconds followed by 60°C for 30 seconds followed by 72°C for 1 minute.

**[0209]** Then, 20  $\mu$ l of the RT-PCR product was subjected to an agarose gel electrophoresis, stained with ethidium

bromide, and irradiated with UV to identify the signals amplified specifically. The gel photograph is shown in Figure 1. As evident from Figure 1, Gm1 is expressed highly in brain, thymus, testis, spleen, small intestine, uterus and heart.

(Example 3) In situ hybridization in brain tissue (Detailed analysis of expression profile of nucleic acid encoding inventive protein in brain tissue)

**[0210]** The expression profile of a nucleic acid encoding the inventive protein in brain tissue was investigated by the following method.

**[0211]** For conducting an in situ hybridization in a mouse brain tissue, the following procedure was employed to clone a cDNA in the 5' terminal region of a mouse Gm1 gene.

**[0212]** 20 ng of a mouse brain-derived cDNA (Clontech) was employed as a template together with 10  $\mu$ M of a forward primer prmGm1-1 (5'-ATGGGCCTATGCTACAGCCTGCGGCCGCT; SEQ ID No:15) and 10  $\mu$ M of a reverse primer prmGm1-2 (5'-GCTGCAGGTCCCGCTTCTGCTCGCGCAGCATGCGGT; SEQ ID No:16) as well as TAKARA LA Taq polymerase (TAKARA LA Taq with GC Buffer, Takara) to perform a PCR to obtain an amplified DNA.

**[0213]** The PCR condition involved 35 cycles, each cycle involving incubations at 95°C for 30 seconds followed by 60°C for 30 seconds followed by 72°C for 2 minutes. The resultant DNA was subjected to an agarose gel electrophoresis followed by a purification with a QIAquick Gel Extraction kit (QIAGEN), and then recovered. This purified and recovered DNA was used as an insert DNA.

**[0214]** Then, a QIAGEN PCR Cloning kit (QIAGEN) was used following to its attached protocol to insert the insert DNA (50ng) into a cloning site of a pDrive vector (10ng), whereby producing a pDrmGm1.

Similarly, a QIAGEN PCR Cloning kit (QIAGEN) was used following to its attached protocol to insert the insert DNA (50ng) to a cloning site of a pDrive vector (10ng), whereby obtaining a pDrmGolf.

(Production of probe for in situ hybridization)

**[0215]** 1  $\mu$ g of a pDrmGm1 plasmid was cleaved with a restriction enzyme HindIII or BamHI to obtain a linear plasmid pDrmGm1/HindIII and pDrmGm1/BamHI. 1  $\mu$ g of a pDrmGm1/HindIII, 2  $\mu$ l of a DIGRNALabelingMix (Roche, Diagnostic) and 1  $\mu$ l of a SP6RNA polymerase (Roche, Diagnostic) were mixed and incubated at 37°C for 3 hours in the presence of the attached buffer. Then, 1  $\mu$ l of a DNaseI (Roche, Diagnostic) was added and the mixture was incubated at 37°C for 30 minutes to obtain a cRNA. This cRNA was precipitated with ethanol, suspended in 20  $\mu$ l of a TE buffer, and used as an mGm1 sense cRNA.

**[0216]** Similarly, 1  $\mu$ g of a pDrmGm1/BamHI, 2  $\mu$ l of a DIGRNALabelingMix (Roche, Diagnostic) and 1  $\mu$ l of SP6RNA polymerase (Roche, Diagnostic) were mixed, and incubated at 37°C for 3 hours in the presence of the attached buffer. Then, 1  $\mu$ l of a DNaseI (Roche, Diagnostic) was added and incubated at 37°C for 30 minutes to obtain a cRNA. This cRNA was precipitated with ethanol, suspended in a 20  $\mu$ l of TE buffer, and used as an mGm1 antisense cRNA.

(Detection by in situ hybridization)

**[0217]** A detailed analysis of the expression profile of mRNA of the Gm1 in a brain is conducted by an in situ hybridization using a labeled cRNA [Simmons et al., J. Histotechnol. 12:169-181(1989)]. Thus, from a mouse brain fixed using paraformaldehyde and glutaraldehyde by a known method, a brain section whose thickness is 50  $\mu$ m is prepared using a brain section producing device (sliding microtome), and then adsorbed on a glass slide and dried. The brain section is made free from the paraffin, autoclaved in a target solution (Daco) (105°C, 10 minutes), dehydrated, and dried in the air. The hybridization with a probe (100ng cRNA) is conducted in a hybridization buffer (40% formamide, 4 $\times$ SSC, 1mMEDTA, 250  $\mu$ g/ml yeast tRNA, 1 $\times$ Denhardt's solution, 10% dextran sulfate) at 60°C overnight. Thereafter, the brain section is washed at 65°C with 2 $\times$ SSC, 0.1% SDS solution, treated with an RNaseA (10  $\mu$ g/ml, 37°C, 30 minutes), washed with 2 $\times$ SSC, 50% formamide solution, dehydrated, dried in the air, and subjected to a mRNA detection using a DIG labeled antibody detection kit (Daco).

(Example 4) Construction of expression plasmid for expression of human Gm1 protein in E.coli

**[0218]** In order to express a large amount of a human Gm1 protein in E.coli, a human Gm1 protein is first expressed as a fusion protein with a glutathion S transferase, and then only the part of the human Gm1 protein is cut out from the fusion protein.

**[0219]** Thus, the human Gm1 cDNA fragment-containing plasmid pCR-Gm1 obtained in Example 1 is double-digested with EcoRV and SpeI, and imparted with a blunt end with a Blunting Kit (Takara). The resultant DNA is subjected to an agarose gel electrophoresis and then purified using a QIAquick Gel Extraction Kit (QIAGEN) and then recovered. The recovered DNA is used as an insert DNA. The pGEX-5X-1 which had been cleaved with EcoRV and then BAP-

treated is employed as a vector, and 50 ng of this vector and 10 ng of the insert DNA are ligated using a T4 ligase, whereby producing an expression plasmid pGEX-Gm1.

(Example 5) Purification of recombinant human Gm1 protein from E.coli expressing glutathion S transferase - human Gm1 fusion protein

**[0220]** The glutathion S transferase - human Gm1 fusion protein-expressing plasmid pGEX-Gm1 obtained in Example 4 is used to transform an E.coli (*Escherichia coli*) JM109 strain by a calcium method. The resultant transformant is cultured in a 50 µg/ml ampicillin (Sigma)-supplemented LB medium at 37°C, and, once the O.D.<sub>600</sub> becomes about 0.6 reached, 1 mM (final concentration) of isopropyl-β-D-thiogalactopyranoside (IPTG) is added to induce the protein expression, and incubated for further 6 hours, prior to the recovery of the cells.

**[0221]** The cells are disrupted with ultrasonic treatment, centrifuged at 10,000g for 5 minutes to obtain a soluble fraction. The resultant soluble fraction is applied onto an anti-glutathion S transferase monoclonal antibody column (Amersham Bioscience) to purify a glutathion S transferase - human Gm1 fusion protein. Then, the purified glutathion S transferase - human Gm1 fusion protein is treated with an active blood coagulation factor X (New England Biolabo) to cut a human Gm1 protein out.

**[0222]** The human Gm1 protein thus cut out is subjected sequentially to a cation exchange column (S-sepharose FF; Pharmacia), hydrophobic column (Phenyl-superose; Pharmacia), hydroxyapatite column (MITSUI TOATSU CHEMICALS), cation exchange column (MONOS; Pharmacia) to purify the human Gm1 protein until it shows an almost single band in an SDS-PAGE analysis with Coomassie brilliant blue staining.

(Example 6) Production of human Gm1 protein partial peptide and production of anti-human Gm1 peptide antibody using this peptide

**[0223]** An antibody specific to a human Gm1 protein was prepared by the procedure shown below. A peptide consisting of 14 amino acids of the amino acid Nos.7 to 20 in the amino acid sequence represented by SEQ ID No:1 was synthesized.

**[0224]** This peptide was bound to a carrier protein KML and used as an immunogen. The resultant KML fusion peptide ptGm1 was used to immunize a New Zealand white rabbit to produce an anti-human Gm1 peptide serum. The immunization was repeated 5 times. From this rabbit, an antiserum was collected, and the antiserum was purified using a protein G column (Amersham Bioscience) to isolate an antigen-specific anti-human Gm1 protein antibody.

(Example 7) Construction of expression vector for expression of human Gm1 protein in animal cell

**[0225]** An expression vector for transient expression of a human Gm1 protein in an animal cell is constructed. Thus, first, the human Gm1 cDNA fragment-containing pCR-Gm1 obtained in Example 1 is double-digested with restriction enzymes XbaI and KpnI, and the resultant DNA fragment is introduced into a pcDNA3.1 at XbaI site and KpnI site whereby obtaining an expression vector pcDNA-Gm1 for a transient expression of human Gm1 protein in an animal cell.

(Example 8) Construction of expression vectors for expression of human dopamine receptor proteins in animal cell

**[0226]** Expression vectors for transient expression in an animal cell of a human dopamine D1 receptor protein and a human dopamine D2 receptor protein respectively were constructed by the following procedure.

**[0227]** In order to amplify a DNA encoding a human dopamine D1 receptor, 20 ng of a plasmid DNA from a human brain-derived cDNA library (Takara) (pAP3neo) was employed as a template together with 10 µM of a forward primer prDopaminD1-5' (5'-agctcggatccATGAGGACTCTGAACACCTCTGCCA; (SEQ ID NO:17) and 10 µM of a reverse primer prDopaminD1-3' (5'-gtgcagaattcTCATCTGCGAGTTCAGGTTGGGT; SEQ ID No:18) as well as a TAKARA LA Taq polymerase (TAKARA LA Taq with GC Buffer, Takara) to perform a PCR.

**[0228]** In order to amplify a DNA encoding a human dopamine D2 receptor, 20 ng of a plasmid DNA from a human brain-derived cDNA library (Takara) (pAP3neo) was used as a template together with 10 µM of a forward primer prDopaminD2-5'

(5' -agctcggatccATGGATCCACTGAATCTGTCCTGGTATGA; SEQ ID No:19)

and 10 µM of a reverse primer prDopaminD2-3' (5'-gtgcagaattcTCAGCAGTGAAGGATCTTCTGGAAGGCCTT; SEQ ID No:20) as well as a TAKARA LA Taq polymerase (TAKARA LA Taq with GC Buffer, Takara) to perform a PCR.

**[0229]** The PCR condition involved 35 cycles, each cycle involving incubations at 95°C for 30 seconds followed by

60°C for 30 seconds followed by 72°C for 2 minutes. The resultant DNA was subjected to an agarose gel electrophoresis followed by a purification with a QIAquick Gel Extraction kit (QIAGEN), and then recovered. This purified and recovered DNA was used as an insert DNA.

[0230] Each insert DNA was double-digested with BamHI and EcoRI, introduced into a pcDNA3.1(+) at BamHI and EcoRI sites, whereby obtaining animal cell expression vectors pcDNA-D1R and pcDNA-D2R.

(Example 9) Construction of Baculovirus vector encoding human Gm1 protein

[0231] First, a transfer vector for overexpressing a human Gm1 protein in an insect cell was constructed. The human Gm1 cDNA fragment-containing pCR-Gm1 obtained in Example 1 was double-digested with restriction enzyme XbaI and SpeI, and the resultant DNA fragment was introduced into a pAcMP2 (Pharmingen) at XbaI site to obtain a transfer vector pAcMP-Gm1. Then 5 µg of this transfer vector and 1 µg of a Baculovirus DNA, BaculoGold DNA (Pharmingen) were cotransfected to 2 x 10<sup>6</sup> cells of Sf21, which were cultured at 27°C for 5 days, and then the culture supernatant was recovered to obtain a virus solution.

(Example 10) Construction of Baculovirus vector encoding human Gβ protein

[0232] A transfer vector for overexpressing a human Gβ protein in an insect cell was constructed by the procedure described below.

[0233] In order to amplify a DNA encoding human Gβ protein, 20 ng of a plasmid DNA from a human brain-derived cDNA library (Takara) (pAP3neo) was employed as a template together with 10 µM of a forward primer prGb1-5' (5'-AT-GAGTGAGCTTGACCAGTTACGGCA; SEQ ID No:21), 10 µM of a reverse primer prGb1-3' (5'-TTAGTTCAGATCTT-GAGGAAGCTAT; SEQ ID No:22) as well as a TAKARA LA Taq polymerase (TAKARA LA Taq with GC Buffer, Takara) to perform a PCR. The PCR condition involved 35 cycles, each cycle involving incubations at 95°C for 30 seconds followed by 60°C for 30 seconds followed by 72°C for 2 minutes. The resultant DNA was subjected to an agarose gel electrophoresis followed by a purification with a QIAquick Gel Extraction kit (QIAGEN), and then recovered. This purified and recovered DNA was used as an insert DNA. Then, a TOPO TA Cloning Kit (Invitrogen) was used in accordance with its attached protocol to insert the resultant insert DNA (50ng) into a pCR2.1-TOPO vector (10ng at a cloning site whereby obtaining a pCR-Gβ.

[0234] Then, the pCR-Gβ was double-digested with restriction enzyme BamHI and NotI, and the resultant DNA fragment was introduced into a pAcMP3 (Pharmingen) at BamHI and NotI sites, whereby obtaining a transfer vector animal cell expression vector pAcMP-Gβ.

Then, 5 µg of this transfer vector and 1 µg of a Baculovirus DNA, BaculoGold DNA (Pharmingen) were cotransfected to 2 x 10<sup>6</sup> cells of Sf21 cell, which are cultured at 27 µ for 5 days, and then the culture supernatant was recovered to obtain a virus solution.

(Example 11) Construction of Baculovirus vector encoding human Gy protein

[0235] A transfer vector for overexpressing a human Gy protein in an animal cell was constructed by the procedure described below.

[0236] In order to amplify a DNA encoding human Gy protein, 20 ng of a plasmid DNA from a human brain-derived cDNA library (Takara) (pAP3neo) was employed as a template together with 10 µM of a forward primer prGg3-5' (5'-AT-GAAAGGTGAGACCCCGGTGAACA; SEQ ID No:23), 10 µM a reverse primer prGg3-3' (5'-TCAGAGGAGAGCACA-GAAGAAGCTT; SEQ ID No:24) as well as a TAKARA LA Taq polymerase (TAKARA LA Taq with GC Buffer, Takara) to perform a PCR. The PCR condition involved 35 cycles, each cycle involving incubations at 95°C for 30 seconds followed by 60°C for 30 seconds followed by 72°C for 2 minutes. The resultant DNA was subjected to an agarose gel electrophoresis followed by a purification with a QIAquick Gel Extraction kit (QIAGEN), and then recovered. This purified and recovered DNA was used as an insert DNA. Then, a TOPO TA Cloning Kit (Invitrogen) was used in accordance with its attached protocol to insert each resultant insert DNA (50ng) into a pCR2.1-TOPO vector (10ng) at a cloning site whereby obtaining a pCR-Gy.

[0237] Then, the pCR-Gy was double-digested with restriction enzyme XbaI and PstI, and the resultant DNA fragment was introduced into a pAcMP3 (Pharmingen) at XbaI and PstI sites, whereby obtaining a transfer vector pAcMP-Gy. Then, 5 µg of this transfer vector and 1 µg of a Baculovirus DNA, BaculoGold DNA (Pharmingen) were cotransfected to 2 x 10<sup>6</sup> cells of Sf21 cell, which are cultured at 27°C for 5 days, and then the culture supernatant was recovered to obtain a virus solution.

(Example 12) Construction of Baculovirus vectors encoding human dopamine D1 receptor and human dopamine D2 receptor

[0238] Transfer vectors for overexpressing a human dopamine D1 receptor protein and a human dopamine D2 receptor protein respectively in an insect cell were constructed by the procedure described below.

[0239] The human dopamine D1 receptor expression vector pcDNA-D1R and the human dopamine D2 receptor expression vector pcDNA-D2R obtained in Example 8 were each double-digested with BamHI and EcoRI, and the resultant DNA fragments are each introduced into a pAcMP3 at BamHI and EcoRI sites, whereby obtaining transfer vectors pAcMP-D1R and pAcMP-D2R.

Then, each 5 µg of the either transfer vectors and 1 µg of a Baculovirus DNA, BaculoGold DNA (Pharmingen) were cotransfected to  $2 \times 10^6$  cells of Sf21 cell, which are cultured at 27 °C for 5 days, and then the culture supernatants were recovered to obtain virus solutions.

(Example 13) High expression of human Gm1 protein using Baculovirus vector and purification of the protein

[0240] A virus solution containing human Gm1 protein expression Baculovirus obtained in Example 9 was infected at MOI5 to  $2 \times 10^7$  cells of SF21 cell, which were cultured at 27°C. Five days after the infection, the cells were recovered and suspended in an HE/PI buffer (20mM HEPSE, 2mM EDTA supplemented with 1Xprotease inhibitor cocktail (NACALAITESQUE)). The cell suspension was passed through a 26G needle 15 times to disrupt the cell membrane. The suspension was then centrifuged at 4°C and 100xg for 5 minutes, and the supernatant obtained was centrifuged at 4°C and 20,000xg for 30 minutes, whereby recovering a human Gm1 protein-containing cell membrane fraction.

(Example 14) High expression of human Gβ protein and human Gy protein using Baculovirus vector and purification of the proteins

[0241] A virus solution containing human Gβ protein expression Baculovirus obtained in Example 10 and human Gy protein expression Baculovirus obtained in Example 11 was infected at MOI5 to  $2 \times 10^7$  cells of SF21 cell, which were cultured at 27°C. Five days after the infection, the cells were recovered and suspended in an HE/PI buffer (20mM HEPSE, 2mM EDTA supplemented with 1XProtease inhibitor cocktail (NACALAITESQUE)). The cell suspension was passed through a 26G needle 15 times to disrupt the cell membrane. The cell suspension was then centrifuged at 4°C and 110xg for 5 minutes, and the supernatant obtained was centrifuged at 4°C and 20,000xg for 30 minutes, whereby recovering a cell membrane fraction containing the human Gβ protein and the human Gy protein.

(Example 15) High expression of human dopamine D1 receptor protein and human dopamine D2 receptor Gy protein using Baculovirus vector and purification of the proteins

[0242] A virus solution containing either human dopamine D1 receptor protein expression Baculovirus or human dopamine D2 receptor protein expression Baculovirus obtained in Example 12 was infected at MOI5 to  $2 \times 10^7$  cells of SF21 cell, which were cultured at 27°C. Five days after the infection, the cells were recovered and suspended in an HE/PI buffer (20mM HEPSE, 2mM EDTA supplemented with 1XProtease inhibitor cocktail (NACALAITESQUE)). The cell suspensions were passed through a 26G needle 15 times to disrupt the cell membrane. This cell suspensions were then centrifuged at 4°C and 110xg for 5 minutes, and the supernatants obtained were centrifuged at 4°C and 20,000xg for 30 minutes, whereby recovering a cell membrane fraction containing the human dopamine D1 receptor protein or the human dopamine D2 receptor protein.

(Example 16) GTP binding assay using human Gm1 protein expressed by Baculovirus vector

[0243] The Gm1 protein-containing membrane fraction purified in Example 13 is employed to conduct a GTP binding assay.

[0244] The cell membrane fraction containing 2 µg of the Gm1 protein prepared in Example 13, the cell membrane fraction containing 2 µg of the Gβ protein and the Gy protein prepared in Example 14 and the cell membrane fraction containing 2 µg of the dopamine D1 receptor protein prepared in Example 15 are suspended in 55 µl of a binding buffer (59mM Tris, 4.8mM MgCl<sub>2</sub>, 2mM EDTA, 100mM NaCl, 1µM GDP). One µM of dopamine is added and the mixture is incubated at 30°C for 10 minutes. Thereafter, 200 pM of [35S] GTPγS is added and the mixture is incubated at 30°C for 30 minutes.

[0245] Then 1.5 ml of a washing buffer (ice-cooled 50mM Tris, 5mM MgCl<sub>2</sub>, 150mM NaCl, 0.1%BSA, 0.05%CHAPS (pH7.4)) is added and the mixture is filtered through a glass fiber filter paper GF/F. Then this filter paper is washed three times with 1 ml of Tris (pH7.4), incubated at 65°C for 30 minutes, subjected to a liquid scintillation counter to

measure the radioactivity of the [35S]GTP $\gamma$ S which is bound to the membrane fraction depositing on the filter paper.

(Example 17) Screening for dopamine D1 receptor antagonist using change in cAMP level as index

5 [0246]  $2 \times 10^5$  Cells of CHO cell were transfected with 1  $\mu$ g of the dopamine D1 receptor expression vector obtained in Example 8 and the Gm1 expression vector obtained in Example 7 (pcDNA-Gm1; 3  $\mu$ g) by a lipofection method to prepare a test cell.

[0247] Then, the cells were inoculated to each well of a 96-well plate at  $3 \times 10^4$  cells/well, and cultured for about 24 hours. Then, the culture medium was removed, and 80  $\mu$ l of 1mM IBMX-supplemented OPTI-MEN (Invitrogen) was added to the cells, which were then incubated at 37°C for 10 minutes.

10 [0248] Then, 10  $\mu$ M of dopamine as a GPCR ligand and 10  $\mu$ M of each test substance (butaclamol, chlorpromazine, fluphenazine, haloperidol, SCH-23390) were added and incubated at 37°C for 30 minutes.

[0249] Then, the reaction buffer was removed, and the cAMP level was determined using a HitHunter ECF cyclic AMP chemiluminescent assay kit (Applied Biosystems). As controls, a test cell which had been brought into contact only with the GPCR ligand at the same concentration and a test cell which had been brought into contact with nothing were examined for their cAMP levels in the similar manner.

15 [0250] A substance which gave a cAMP level percentage of 85% or less upon contact with the ligand and the test substance was selected as a signal transduction inhibitor (antagonist), on the basis of the cAMP level with no contact being 0% and the cAMP level with the contact only with the ligand being 100% (Figure 2).

20 (Example 18) Screening of dopamine D1 receptor using change in cAMP level as index

[0251]  $2 \times 10^5$  Cells of CHO cell were transfected with the dopamine D1 receptor expression vector (1  $\mu$ g) obtained in Example 8 and the Gm1 expression vector obtained in Example 7 (pcDNA-Gm1; 3  $\mu$ g) by a lipofection method to prepare a test cell.

25 [0252] Then, the cells were inoculated to each well of a 96-well plate at  $3 \times 10^4$  cells/well, and cultured for about 24 hours. Then, the culture medium was removed, and 80  $\mu$ l of 1mM IBMX-supplemented OPTI-MEN (Invitrogen) was added to the cells, which were then incubated at 37°C for 10 minutes.

[0253] Then, 10  $\mu$ M of dopamine as a GPCR ligand or 10  $\mu$ M of a test substance (apomorphine, CY208-248, SKF-36393, SKF-81297) was added and incubated at 37°C for 30 minutes.

30 [0254] Then, the reaction buffer was removed, and the cAMP level was determined using a HitHunter ECF cyclic AMP chemiluminescent assay kit (Applied Biosystems). As a control, a test cell which had been brought into contact with nothing were examined for their cAMP levels in the similar manner.

35 [0255] A substance which gave a cAMP level percentage of 125% or more upon contact of the test cell with the test substance was selected as a signal transduction activator (agonist), on the basis of the cAMP level when the test cell has not been brought into contact with anything being 100% (Figure 3).

(Example 19) Screening using change in cAMP level as index

40 [0256]  $1 \times 10^6$  Cells of CHO cell are transfected with the dopamine D1 receptor expression vector obtained in Example 8 (100 ng), CRE-reporter plasmid (pCRE-1uc; 20ng; Stratagene) and the Gm1 expression vector obtained in Example 7 (pcDNA-Gm1; 30 ng) by a lipofection method to prepare a test cell.

[0257] Then the cells are inoculated to each well of a 24-well plate at  $5 \times 10^3$  cells/well, and then cultured for about 48 hours. Then, the cells are washed with 0.2mM buffer (3-isobutyl-methylxanthine, 0.05% BSA, 20mM HEPES-supplemented Hunk's buffer (pH7.4); hereinafter referred to as "reaction buffer"). Then the reaction buffer is added to the cells, which are incubated at 37°C for 30 minutes.

45 [0258] Then, the reaction buffer is removed, and 0.25 ml of a fresh reaction buffer is added to the cells, and then 1 nM dopamine as a GPCR ligand and 0.1 nM to 10 nM test substance are added and the mixture is incubated at 37°C for 30 minutes. Then the cells are dissolved in a cell lysis solution (PICK-A-GENE luciferase kit, TOYO INK), and combined with a luminescent substrate (PICK-A-GENE luciferase kit, TOYO INK), and examined for the fluorescent intensity using a luminometer. As controls, a test cell which had been brought into contact only with the GPCR ligand at the same concentration and a test cell which had been brought into contact with nothing are examined for their fluorescent intensities in the similar manner.

50 [0259] A substance which gave a fluorescence intensity of percentage of 50% or less, or 150% or more upon contact with the ligand and the test substance is selected as a signal transduction regulating substance, on the basis of the fluorescent intensity with no contact being 0% and the fluorescent intensity with the contact only with the ligand being 100%.

(Example 20) Construction of expression vector for expression of human adenosine A2a receptor protein in animal cell

**[0260]** An expression vector for transient expression of a human adenosine A2a receptor protein in an animal cell was constructed by the procedure described below.

**[0261]** In order to amplify a DNA encoding a human adenosine A2a receptor, 20 ng of a plasmid DNA from a human brain-derived cDNA library (Takara) (pAP3neo) was employed as a template together with 10  $\mu$ M of a forward primer prAdenosineA2A-5' (5'-agctcggatccATGCCCATCATGGGCTCCTCGGTGTA; SEQ ID NO:33) and 10  $\mu$ M of a reverse primer prAdenosineA2A-3' (5'-gtgcagaattcTCAGGACACTCCTGCTCCATCCT; SEQ ID No:34) as well as a TAKARA LA Taq polymerase (TAKARA LA Taq with GC Buffer, Takara) to perform a PCR.

**[0262]** The PCR condition involved 35 cycles, each cycle involving incubations at 95°C for 30 seconds followed by 60°C for 30 seconds followed by 72°C for 2 minutes. The resultant DNA was subjected to an agarose gel electrophoresis followed by a purification with a QIAquick Gel Extraction kit (QIAGEN), and then recovered. This purified and recovered DNA was used as an insert DNA.

**[0263]** Each insert DNA was double-digested with BamHI and EcoRI, introduced into a pcDNA3.1(+) at BamHI and EcoRI sites, whereby obtaining an animal cell expression vector pcDNA-A2a.

(Example 21) Screening of adenosine A2a receptor antagonist using change in cAMP level as index

**[0264]**  $2 \times 10^5$  Cells of CHO cell were transfected with the adenosine A2a receptor expression vector (1  $\mu$ g) obtained in Example 20 and the Gm1 expression vector obtained in Example 7 (pcDNA-Gm1; 3  $\mu$ g) by a lipofection method to prepare a test cell.

**[0265]** Then, the cells were inoculated to each well of a 96-well plate at  $3 \times 10^4$  cells/well, and cultured for about 24 hours. Then, the culture medium was removed, and 80  $\mu$ l of 1mM IBMX-supplemented OPTI-MEN (Invitrogen) was added to the cells, which were then incubated at 37°C for 10 minutes.

**[0266]** Then, 10  $\mu$ M of adenosine as a GPCR ligand and 10  $\mu$ M of a test substance (DMPX) were added and incubated at 37°C for 30 minutes.

**[0267]** Then, the reaction buffer was removed, and the cAMP level was determined using a HitHunter ECF cyclic AMP chemiluminescent assay kit (Applied Biosystems). As controls, a test cell which had been brought into contact only with the GPCR ligand at the same concentration and a test cell which had been brought into contact with nothing were examined for their cAMP levels in the similar manner.

**[0268]** A substance which gave a cAMP level percentage of 85% or less upon contact of the cell with the ligand and the test substance was selected as a signal transduction inhibitor (antagonist), on the basis of the cAMP level with no contact being 0% and the cAMP level with the contact of the cell only with the ligand being 100% (Figure 4).

(Example 22) Screening of adenosine A2a receptor agonist using change in cAMP level as index

**[0269]**  $2 \times 10^5$  Cells of CHO cell were transfected with the adenosine A2a receptor expression vector (1  $\mu$ g) obtained in Example 20 and the Gm1 expression vector obtained in Example 7 (pcDNA-Gm1; 3  $\mu$ g) by a lipofection method to prepare a test cell.

**[0270]** Then, the cells were inoculated to each well of a 96-well plate at  $3 \times 10^4$  cells/well, and cultured for about 24 hours. Then, the culture medium was removed, and 80  $\mu$ l of 1mM IBMX-supplemented OPTI-MEN (Invitrogen) was added to the cells, which were then incubated at 37°C for 10 minutes.

**[0271]** Then, 10  $\mu$ M of adenosine as a GPCR ligand or 10  $\mu$ M of a test substance (CGS-21680) was added and incubated at 37°C for 30 minutes.

**[0272]** Then, the reaction buffer was removed, and the cAMP level was determined using a HitHunter ECF cyclic AMP chemiluminescent assay kit (Applied Biosystems). As a control, a test cell which had been brought into contact with nothing were examined for their cAMP levels in the similar manner.

**[0273]** A substance which gave a cAMP level percentage of 125% or more upon contact with the test substance was selected as a signal transduction activator (agonist), on the basis of the cAMP level when the test cell has not been brought into contact with anything being 100% (Figure 5).

(Example 23) Screening by GTP binding assay

**[0274]** The cell membrane fraction containing 2  $\mu$ g of the Gm1 protein prepared in Example 13, the cell membrane fraction containing 2  $\mu$ g of the G $\beta$  protein and G $\gamma$  protein prepared in Example 14 and the cell membrane fraction containing 2  $\mu$ g of the dopamine D1 receptor protein prepared in Example 15 are suspended in 55  $\mu$ l of a binding buffer (59mM Tris, 4.8mM MgCl<sub>2</sub>, 2mM EDTA, 100mM NaCl, 1  $\mu$ M GDP). 1  $\mu$ M of dopamine and 0.1 nM - 10 nM of a test substance (for example, 1 nM SCH-23390) are added and the mixture is incubated at 30°C for 10 minutes. Thereafter,



200 pM of [35S]GTP $\gamma$ S is added and the mixture is incubated at 30°C for 30 minutes.

**[0275]** Then 1.5 ml of a washing buffer (ice-cooled 50mM Tris, 5mM MgCl<sub>2</sub>, 150mM NaCl, 0.1%BSA, 0.05%CHAPS (pH7.4)) is added and the mixture is filtered through a glass fiber filter paper GF/F. Then this filter paper is washed three times with 1 ml of Tris (pH7.4), incubated at 65°C for 30 minutes, subjected to a liquid scintillation counter to

measure the radioactivity of the [35S]GTP $\gamma$ S which is bound to the membrane fraction depositing on the filter paper.  
**[0276]** A substance whose radioactivity percentage is calculated to be 50% or less, or 150% or more upon contact with the both of the ligand and the test substance is selected as a substance capable of regulating the signal transduction, on the basis of the radioactivity with the addition only of the ligand being 100% and the radioactivity without the addition of the ligand or the test substance being 0%.

(Example 24) Assay of activation of signal transduction pathway mediated by Gm1 using change in cAMP level as index

**[0277]** 2 x 10<sup>5</sup> Cells of CHO cell were transfected with the dopamine D1 receptor expression vector (1  $\mu$ g) obtained in Example 8 and the Gm1 expression vector obtained in Example 7 (pcDNA-Gm1; 3  $\mu$ g) by a lipofection method to prepare a test cell.

**[0278]** Then, the cells were inoculated to each well of a 96-well plate at 3 x 10<sup>4</sup> cells/well, and cultured for about 24 hours. Then, the culture medium was removed, and 80  $\mu$ l of 1mM IBMX-supplemented OPTI-MEN (Invitrogen) was added to the cells, which were then incubated at 37°C for 10 minutes.

**[0279]** Then, 10  $\mu$ M of dopamine as a GPCR ligand was added and incubated at 37°C for 30 minutes.

**[0280]** Then, the reaction buffer was removed, and the cAMP level was determined using a HitHunter ECF cyclic AMP chemiluminescent assay kit (Applied Biosystems). As a control, a test cell which had been brought into contact with nothing were examined for their cAMP levels in the similar manner.

**[0281]** The cAMP level of the test cell expressing the Gm1 and the dopamine D1 receptor was determined on the basis of the cAMP level upon no contact being regarded to be 0% while the cAMP level of the control cell expressing only the dopamine D1 receptor being regarded to be 100%, and was revealed to be 127%. Therefore, it was proven that the Gm1-mediated signal transduction system do exist (Figure 6).

(Example 25) Assay of activation of signal transduction pathway mediated by Gm1 using change in cAMP level as index

**[0282]** 2 x 10<sup>5</sup> Cells of CHO cell were transfected with the adenosine A2a receptor expression vector (1  $\mu$ g) obtained in Example 8 and the Gm1 expression vector obtained in Example 7 (pcDNA-Gm1; 3  $\mu$ g) by a lipofection method to prepare a test cell.

**[0283]** Then, the cells were inoculated to each well of a 96-well plate at 3 x 10<sup>4</sup> cells/well, and cultured for about 24 hours. Then, the culture medium was removed, and 80  $\mu$ l of 1mM IBMX-supplemented OPTI-MEN (Invitrogen) was added to the cells, which were then incubated at 37°C for 10 minutes.

**[0284]** Then, 10  $\mu$ M of adenosine as a GPCR ligand was added and incubated at 37°C for 30 minutes.

**[0285]** Then, the reaction buffer was removed, and the cAMP level was determined using a HitHunter ECF cyclic AMP chemiluminescent assay kit (Applied Biosystems). As a control, a test cell which had been brought into contact with nothing were examined for their cAMP levels in the similar manner.

**[0286]** The cAMP level when allowing the Gm1 and the adenosine A2a receptor to be expressed was determined on the basis of the cAMP level upon no contact being regarded to be 0% while the cAMP level of the cell allowed to express only the adenosine A2a receptor being regarded to be 100%, and was revealed to be 134%. Therefore, it was proven that the Gm1-mediated signal transduction system do exist (Figure 7).

(Example 26) Cloning of mouse Gm1 protein-encoding cDNA

**[0287]** A pDr-mGm1 which is a plasmid having a DNA encoding the full-length mouse Gm1 was produced as described below.

**[0288]** 20 ng of a mouse brain-derived cDNA library (Clontech) was employed as a template together with 10  $\mu$ M of a forward primer (5'-ATGGGCCTATGCTACAGCCTGCGGCCGCT; SEQ ID No:29) and 10  $\mu$ M of a reverse primer prmGm1STOP (5'-TCACAAGAGTTCGTAAGTCTGCTTGGATGCTTCT; SEQ ID No:30) as well as a TAKARA LA Taq polymerase (TAKARA LA Taq with GC Buffer, Takara) to conduct a PCR to obtain an amplified DNA.

**[0289]** The PCR condition involved 35 cycles, each cycle involving incubations at 95°C for 30 seconds followed by 60°C for 30 seconds followed by 72°C for 2 minutes. The resultant DNA was subjected to an agarose gel electrophoresis followed by a purification with a QIAquick Gel Extraction kit (QIAGEN), and then recovered. This purified and recovered DNA was used as an insert DNA.

**[0290]** Then, a QIAGEN PCR Cloning kit (QIAGEN) was used following to its attached protocol to insert the insert DNA (50 ng) into a cloning site of a pDrive vector (25 ng), whereby producing a pDr-mGm1.

[0291] The DNA thus obtained was subjected to an ABI377DNA sequencer to determine the nucleotide sequence, which was revealed to contain the nucleotide Nos.1 to 1347 in the nucleotide sequence represented by SEQ ID No: 27 and to encode the full-length amino acid sequence represented by SEQ ID No:25.

5 (Example 27) Cloning of rat Gm1 protein-encoding cDNA

[0292] A pDr-rGm1 which is a plasmid having a DNA encoding the full-length rat Gm1 was produced as described below.

10 [0293] 20 ng of a rat brain-derived cDNA library (Clontech) was employed as a template together with 10  $\mu$ M of a forward primer prrGm1ATG (5'-ATGGGCCTGTGCTACAGCCTACGGCCGCTG; SEQ ID No:31) and 10  $\mu$ M of a reverse primer prrGm1'STOP (5'-TCACAAGAGTTCGTACTGCTTGAGGTGCATTCT; SEQ ID No:32) as well as a TAKARA LA Taq polymerase (TAKARA LA Taq with GC Buffer, Takara) to conduct a PCR to obtain an amplified DNA.

15 [0294] The PCR condition involved 35 cycles, each cycle involving incubations at 95°C for 30 seconds followed by 60°C for 30 seconds followed by 72°C for 2 minutes. The resultant DNA was subjected to an agarose gel electrophoresis followed by a purification with a QIAquick Gel Extraction kit (QIAGEN), and then recovered. This purified and recovered DNA was used as an insert DNA.

[0295] Then, a QIAGEN PCR Cloning kit (QIAGEN) was used following to its attached protocol to insert the insert DNA (50 ng) into a cloning site of a pDrive vector (25 ng), whereby producing a pDr-rGm1.

20 [0296] The DNA thus obtained was subjected to an ABI377DNA sequencer to determine the nucleotide sequence, which was revealed to contain the nucleotide Nos.1 to 1353 in the nucleotide sequence represented by SEQ ID No: 28 and to encode the full-length amino acid sequence represented by SEQ ID No:26.

Free text in Sequence Listing

25 [0297]

SEQ ID No:3

an example of the ribozyme of the present invention

SEQ ID No:4

30 an example of the ribozyme of the present invention

SEQ ID No:5

an example of the ribozyme of the present invention

SEQ ID No:6

an example of the ribozyme of the present invention

35 SEQ ID No:7

an example of the ribozyme of the present invention

SEQ ID No:8

an example of the ribozyme of the present invention

SEQ ID No:9

40 an example of the oligonucleotide of the present invention

SEQ ID No:10

an example of the oligonucleotide of the present invention

SEQ ID No:11

a primer used in an example of the present invention

45 SEQ ID No:12

a primer used in an example of the present invention

SEQ ID No:13

a primer used in an example of the present invention

SEQ ID No:14

50 a primer used in an example of the present invention

SEQ ID No:15

a primer used in an example of the present invention

SEQ ID No:16

a primer used in an example of the present invention

55 SEQ ID No:17

a primer used in an example of the present invention

SEQ ID No:18

a primer used in an example of the present invention

SEQ ID No:19  
a primer used in an example of the present invention  
SEQ ID No:20  
a primer used in an example of the present invention  
5 SEQ ID No:21  
a primer used in an example of the present invention  
SEQ ID No:22  
a primer used in an example of the present invention  
SEQ ID No:23  
10 a primer used in an example of the present invention  
SEQ ID No:24  
a primer used in an example of the present invention  
SEQ ID No:29  
a primer used in an example of the present invention  
15 SEQ ID No:30  
a primer used in an example of the present invention  
SEQ ID No:31  
a primer used in an example of the present invention  
SEQ ID No:32  
20 a primer used in an example of the present invention  
SEQ ID No:33  
a primer used in an example of the present invention  
SEQ ID No:34  
25 a primer used in an example of the present invention

SEQUENCE LISTING

5 <110> Sumitomo Chemical Company Limited

10 <120> NOVEL G PROTEINS, POLYNUCLEOTIDE ENCODING THE SAME AND  
UTILIZATION THEREOF

15 <130>

20 <150> JP 2002/206841

<151> 2002-07-16

25 <150> JP 2002/367778

<151> 2002-12-19

30 <150> JP 2003/095955

35 <151> 2003-03-31

40 <160> 34

<210> 1

45 <211> 458

<212> PRT

50 <213> Homo sapiens

55 <400> 1

EP 1 382 613 A1

5 Met Gly Leu Cys Tyr Ser Leu Arg Pro Leu Leu Phe Gly Gly Pro Gly  
 1 5 10 15  
 Asp Asp Pro Cys Ala Ala Ser Glu Pro Pro Val Glu Asp Ala Gln Pro  
 10 20 25 30  
 Ala Pro Ala Pro Ala Leu Ala Pro Val Arg Ala Ala Ala Arg Asp Thr  
 15 35 40 45  
 Ala Arg Thr Leu Leu Pro Arg Gly Gly Glu Gly Ser Pro Ala Cys Ala  
 50 55 60  
 20 Arg Pro Lys Ala Asp Lys Pro Lys Glu Lys Arg Gln Arg Thr Glu Gln  
 65 70 75 80  
 25 Leu Ser Ala Glu Glu Arg Glu Ala Ala Lys Glu Arg Glu Ala Val Lys  
 85 90 95  
 30 Glu Ala Arg Lys Val Ser Arg Gly Ile Asp Arg Met Leu Arg Asp Gln  
 100 105 110  
 Lys Arg Asp Leu Gln Gln Thr His Arg Leu Leu Leu Leu Gly Ala Gly  
 35 115 120 125  
 Glu Ser Gly Lys Ser Thr Ile Val Lys Gln Met Arg Ile Leu His Val  
 40 130 135 140  
 Asn Gly Phe Asn Pro Glu Glu Lys Lys Gln Lys Ile Leu Asp Ile Arg  
 45 145 150 155 160  
 Lys Asn Val Lys Asp Ala Ile Val Thr Ile Val Ser Ala Met Ser Thr  
 50 165 170 175  
 Ile Ile Pro Pro Val Pro Leu Ala Asn Pro Glu Asn Gln Phe Arg Ser  
 180 185 190  
 55 Asp Tyr Ile Lys Ser Ile Ala Pro Ile Thr Asp Phe Glu Tyr Ser Gln

EP 1 382 613 A1

	195	200	205
5	Glu Phe Phe Asp His Val Lys Lys Leu Trp Asp Asp Glu Gly Val Lys		
	210	215	220
10	Ala Cys Phe Glu Arg Ser Asn Glu Tyr Gln Leu Ile Asp Cys Ala Gln		
	225	230	235 240
15	Tyr Phe Leu Glu Arg Ile Asp Ser Val Ser Leu Val Asp Tyr Thr Pro		
	245	250	255
20	Thr Asp Gln Asp Leu Leu Arg Cys Arg Val Leu Thr Ser Gly Ile Phe		
	260	265	270
25	Glu Thr Arg Phe Gln Val Asp Lys Val Asn Phe His Met Phe Asp Val		
	275	280	285
30	Gly Gly Gln Arg Asp Glu Arg Arg Lys Trp Ile Gln Cys Phe Asn Asp		
	290	295	300
35	Val Thr Ala Ile Ile Tyr Val Ala Ala Cys Ser Ser Tyr Asn Met Val		
	305	310	315 320
40	Ile Arg Glu Asp Asn Asn Thr Asn Arg Leu Arg Glu Ser Leu Asp Leu		
	325	330	335
45	Phe Glu Ser Ile Trp Asn Asn Arg Trp Leu Arg Thr Ile Ser Ile Ile		
	340	345	350
50	Leu Phe Leu Asn Lys Gln Asp Met Leu Ala Glu Lys Val Leu Ala Gly		
	355	360	365
55	Lys Ser Lys Ile Glu Asp Tyr Phe Pro Glu Tyr Ala Asn Tyr Thr Val		
	370	375	380
	Pro Glu Asp Ala Thr Pro Asp Ala Gly Glu Asp Pro Lys Val Thr Arg		
	385	390	395 400

EP 1 382 613 A1

Ala Lys Phe Phe Ile Arg Asp Leu Phe Leu Arg Ile Ser Thr Ala Thr

5 405 410 415

Gly Asp Gly Lys His Tyr Cys Tyr Pro His Phe Thr Cys Ala Val Asp

10 420 425 430

Thr Glu Asn Ile Arg Arg Val Phe Asn Asp Cys Arg Asp Ile Ile Gln

15 435 440 445

Arg Met His Leu Lys Gln Tyr Glu Leu Leu

20 450 455

<210> 2

25 <211> 1377

<212> DNA

30 <213> Homo sapiens

<220>

35 <221> CDS

<222> (1).. (1377)

40 <400> 2

45 atg ggt ctg tgc tac agt ctg cgg ccg ctg ctt ttc ggg ggc cca ggg 48

Met Gly Leu Cys Tyr Ser Leu Arg Pro Leu Leu Phe Gly Gly Pro Gly

50 1 5 10 15

gac gac ccc tgc gcg gcc tgg gag ccg ccg gtg gag gac gcg cag ccc 96

55 Asp Asp Pro Cys Ala Ala Ser Glu Pro Pro Val Glu Asp Ala Gln Pro

EP 1 382 613 A1

	20	25	30	
5	gcc ccg gcc ccg gcc ctg gcc cca gtc cgg gcg gcc gca agg gac acg 144			
	Ala Pro Ala Pro Ala Leu Ala Pro Val Arg Ala Ala Ala Arg Asp Thr			
10	35	40	45	
15	gcc egg acc ctg ctc cct cgg ggc ggc gaa ggg agc ccg gca tgc gct 192			
	Ala Arg Thr Leu Leu Pro Arg Gly Gly Glu Gly Ser Pro Ala Cys Ala			
20	50	55	60	
25	cgg ccc aaa gca gac aag ccg aag gag aag cgg cag cgc acc gag cag 240			
	Arg Pro Lys Ala Asp Lys Pro Lys Glu Lys Arg Gln Arg Thr Glu Gln			
30	65	70	75	80
35	ctg agt gcc gag gag cgc gag gcg gcc aag gag cgc gag gcg gtc aag 288			
	Leu Ser Ala Glu Glu Arg Glu Ala Ala Lys Glu Arg Glu Ala Val Lys			
40	85	90	95	
45	gag gcg agg aaa gtg agc cgg ggc atc gac cgc atg ctg cgc gac cag 336			
	Glu Ala Arg Lys Val Ser Arg Gly Ile Asp Arg Met Leu Arg Asp Gln			
50	100	105	110	
55	aag cgc gac ctg cag cag acg cac cgg ctc ctg ctg ctc ggg gct ggt 384			
	Lys Arg Asp Leu Gln Gln Thr His Arg Leu Leu Leu Leu Gly Ala Gly			
	115	120	125	



EP 1 382 613 A1

5 gag tct ggg aaa agc acc atc gtc aaa cag atg agg atc ctg cac gtc 432  
 Glu Ser Gly Lys Ser Thr Ile Val Lys Gln Met Arg Ile Leu His Val

10 130 135 140

15 aat ggg ttt aat ccc gag gaa aag aaa cag aaa att ctg gac atc cgg 480  
 Asn Gly Phe Asn Pro Glu Glu Lys Lys Gln Lys Ile Leu Asp Ile Arg

20 145 150 155 160

25 aaa aat gtt aaa gat gct atc gtg aca att gtt tca gca atg agt act 528  
 Lys Asn Val Lys Asp Ala Ile Val Thr Ile Val Ser Ala Met Ser Thr

30 165 170 175

35 ata ata cct cca gtt ccg ctg gcc aac cct gaa aac caa ttt cga tca 576  
 Ile Ile Pro Pro Val Pro Leu Ala Asn Pro Glu Asn Gln Phe Arg Ser

40 180 185 190

45 gac tac atc aag agc ata gcc cct atc act gac ttt gaa tat tcc cag 624  
 Asp Tyr Ile Lys Ser Ile Ala Pro Ile Thr Asp Phe Glu Tyr Ser Gln

50 195 200 205

55 gaa ttc ttt gac cat gtg aaa aaa ctt tgg gac gat gaa ggc gtg aag 672  
 Glu Phe Phe Asp His Val Lys Lys Leu Trp Asp Asp Glu Gly Val Lys

210 215 220

EP 1 382 613 A1

5 gca tgc ttt gag aga tcc aac gaa tac cag ctg att gac tgt gca caa 720  
Ala Cys Phe Glu Arg Ser Asn Glu Tyr Gln Leu Ile Asp Cys Ala Gln  
225 230 235 240

10 tac ttc ctg gaa aga atc gac agc gtc agc ttg gtt gac tac aca ccc 768  
Tyr Phe Leu Glu Arg Ile Asp Ser Val Ser Leu Val Asp Tyr Thr Pro  
15 245 250 255

20 aca gac cag gac ctc ctc aga tgc aga gtt ctg aca tct ggg att ttt 816  
Thr Asp Gln Asp Leu Leu Arg Cys Arg Val Leu Thr Ser Gly Ile Phe  
25 260 265 270

30 gag aca cga ttc caa gtg gac aaa gta aac ttc cac atg ttt gat gtt 864  
Glu Thr Arg Phe Gln Val Asp Lys Val Asn Phe His Met Phe Asp Val  
35 275 280 285

40 ggt ggc cag agg gat gag agg aga aaa tgg atc cag tgc ttt aac gat 912  
Gly Gly Gln Arg Asp Glu Arg Arg Lys Trp Ile Gln Cys Phe Asn Asp  
290 295 300

45 gtc aca gct atc att tac gtc gca gcc tgc agt agc tac aac atg gtg 960  
Val Thr Ala Ile Ile Tyr Val Ala Ala Cys Ser Ser Tyr Asn Met Val  
50 305 310 315 320

55 att cga gaa gat aac aac acc aac agg ctg aga gag tcc ctg gat ctt 1008

EP 1 382 613 A1

Ile Arg Glu Asp Asn Asn Thr Asn Arg Leu Arg Glu Ser Leu Asp Leu

5 325 330 335

10 ttt gaa agc atc tgg aac aac agg tgg tta cgg acc att tct atc atc 1056

Phe Glu Ser Ile Trp Asn Asn Arg Trp Leu Arg Thr Ile Ser Ile Ile

15 340 345 350

20 ttg ttc ttg aac aaa caa gat atg ctg gca gaa aaa gtc ttg gca ggg 1104

Leu Phe Leu Asn Lys Gln Asp Met Leu Ala Glu Lys Val Leu Ala Gly

355 360 365

25  
30 aaa tca aaa att gaa gac tat ttc cca gaa tat gca aat tat act gtt 1152

Lys Ser Lys Ile Glu Asp Tyr Phe Pro Glu Tyr Ala Asn Tyr Thr Val

370 375 380

35 cct gaa gac gca aca cca gat gca gga gaa gat ccc aaa gtt aca aga 1200

Pro Glu Asp Ala Thr Pro Asp Ala Gly Glu Asp Pro Lys Val Thr Arg

40 385 390 395 400

45 gcc aag ttc ttt atc cgg gac ctg ttt ttg agg atc agc acg gcc acc 1248

Ala Lys Phe Phe Ile Arg Asp Leu Phe Leu Arg Ile Ser Thr Ala Thr

50 405 410 415

55 ggt gac ggc aaa cat tac tgc tac ccg cac ttc acc tgc gcc gtg gac 1296

Gly Asp Gly Lys His Tyr Cys Tyr Pro His Phe Thr Cys Ala Val Asp

420

425

430

5

aca gag aac atc cgc agg gtg ttc aac gac tgc cgc gac atc atc cag 1344

10

Thr Glu Asn Ile Arg Arg Val Phe Asn Asp Cys Arg Asp Ile Ile Gln

435

440

445

15

cgg atg cac ctc aag cag tat gag ctc ttg tga 1377

Arg Met His Leu Lys Gln Tyr Glu Leu Leu

20

450

455

25

<210> 3

<211> 44

<212> RNA

30

<213> artificial sequence

35

<220>

<223> an example of the ribozyme of the present invention

40

<400> 3

45

tcgcctcctt ctgatgaggc cgaaaggccg aaaccgcctc gcgc 44

50

<210> 4

<211> 44

<212> RNA

55

<213> artificial sequence

5 <220>  
<223> an example of the ribozyme of the present invention

10

<400> 4

15 cggccgcccc ctgatgaggc cgaaaggccg aaactggggc cagc 44

20

<210> 5

<211> 43

<212> RNA

25

<213> artificial sequence

30

<220>

<223> an example of the ribozyme of the present invention

35

<400> 5

cagcggccgc ctgatgaggc cgaaaggccg aaactgtagc aca 43

40

<210> 6

45

<211> 54

<212> RNA

50

<213> artificial sequence

55

<220>

<223> an example of the ribozyme of the present invention

5 <400> 6  
tcgcctcctt agaagcctac cagagaaaca cacgttgttg tatattacct ggta 54

10  
15 <210> 7  
<211> 54  
<212> RNA  
<213> artificial sequence

20  
25 <220>  
<223> an example of the ribozyme of the present invention

30 <400> 7  
cggccgcccc agaaggggac cagagaaaca cacgttgttg tatattacct ggta 54

35 <210> 8  
<211> 55  
40 <212> RNA  
<213> artificial sequence

45  
50 <220>  
<223> an example of the ribozyme of the present invention

55 <400> 8  
cagcggccgc aagaagtaga ccagagaaac acacgttgtg gtatattacc tgta 55

<210> 9

&lt;211&gt; 24

&lt;212&gt; DNA

<213> artificial sequence

 $\langle 220 \rangle$ 

<223> an example of the oligonucleotide of the present invention

&lt;400&gt; 9

atgggtctgt gctacagtct gcgg

24

<210> 10

&lt;211&gt; 37

## <212> DNA

⟨213⟩ artificial sequence

 $\langle 220 \rangle$ 

<223> an example of the oligonucleotide of the present invention

<400> 10

acgatggtgc ttttcccaga ctcaccagcc ccgagca

37

<210> 11

&lt;211&gt; 24

<212> DNA

5 <213> artificial sequence

10 <220>

<223> a primer used in an example of the present invention

15 <400> 11

atgggtctgt gctacagtct gcgg

24

20

<210> 12

25 <211> 21

<212> DNA

30 <213> artificial sequence

<220>

35 <223> a primer used in an example of the present invention

40 <400> 12

tcacaagagc tcatactgct t

21

45

<210> 13

50 <211> 22

<212> DNA

55 <213> artificial sequence



<220>

<223> a primer used in an example of the present invention

<400> 13

atgggggtgtt tgggcggcaa ca

22

<210> 14

<211> 37

<212> DNA

<213> artificial sequence

<220>

<223> a primer used in an example of the present invention

<400> 14

acgatggtgc ttttcccaga ctcaccagcc ccgagca

37

<210> 15

<211> 29

<212> DNA

<213> artificial sequence

<220>

<223> a primer used in an example of the present invention

<400> 15

5 atgggcctat gctacagcct gcggccgct 29

<210> 16

10

<211> 36

<212> DNA

15

<213> artificial sequence

<220>

20

<223> a primer used in an example of the present invention

25

<400> 16

30 gctgcaggtc ccgcttctgc tcgcgcagca tgcggt 36

<210> 17

35

<211> 36

<212> DNA

40

<213> artificial sequence

<220>

45

<223> a primer used in an example of the present invention

50

<400> 17

55 agctcgatc catgaggact ctgaacacct ctgcc 36

<210> 18

<211> 34

<212> DNA

<213> artificial sequence

<220>

<223> a primer used in an example of the present invention

<400> 18

gtgcagaatt ctcacctgcg agttcaggtt gggt

34

<210> 19

<211> 40

<212> DNA

<213> artificial sequence

<220>

<223> a primer used in an example of the present invention

<400> 19

agctcggatc catggatcca ctgaatctgt cctggtatga

40

<210> 20

<211> 41

<212> DNA

<213> artificial sequence

5

<220>

<223> a primer used in an example of the present invention

10

<400> 20

15

gtgcagaatt ctcagcagtg aaggatcttc tggaaggcct t

41

20

<210> 21

<211> 26

25

<212> DNA

<213> artificial sequence

30

<220>

<223> a primer used in an example of the present invention

35

<400> 21

40

atgagtgagc ttgaccagtt acggca

26

45

<210> 22

<211> 26

<212> DNA

50

<213> artificial sequence

55

<220>

<223> a primer used in an example of the present invention

5

<400> 22

10

ttagttccag atcttgagga agctat

26

<210> 23

15

<211> 25

<212> DNA

20

<213> artificial sequence

<220>

25

<223> a primer used in an example of the present invention

30

<400> 23

atgaaagggtg agaccccggt gaaca

25

35

<210> 24

40

<211> 24

<212> DNA

<213> artificial sequence

45

<220>

50

<223> a primer used in an example of the present invention

55

<400> 24

tcagaggaga gcacagaaga actt

24

5

&lt;210&gt; 25

10

&lt;211&gt; 448

&lt;212&gt; PRT

15

&lt;213&gt; Mus musculus

20

&lt;400&gt; 25

Met Gly Leu Cys Tyr Ser Leu Arg Pro Leu Leu Phe Gly Ser Pro Glu

1

5

10

15

25

Asp Thr Pro Cys Ala Ala Ser Glu Pro Cys Ala Glu Asp Ala Gln Pro

20

25

30

30

Ser Ala Ala Pro Ala Pro Ala Ser Ile Pro Ala Pro Ala Pro Val Gly

35

40

45

35

Thr Leu Leu Arg Arg Gly Gly Gly Arg Ile Val Ala Asn Ala Arg Pro

50

55

60

40

Pro Gly Glu Leu Gln Ser Arg Arg Arg Gln Glu Gln Leu Arg Ala Glu

65

70

75

80

Glu Arg Glu Ala Ala Lys Glu Ala Arg Lys Val Ser Arg Gly Ile Asp

45

85

90

95

Arg Met Leu Arg Glu Gln Lys Arg Asp Leu Gln Gln Thr His Arg Leu

100

105

110

50

Leu Leu Leu Gly Ala Gly Glu Ser Gly Lys Ser Thr Ile Val Lys Gln

115

120

125

55

Met Arg Ile Leu His Val Asn Gly Phe Asn Pro Glu Glu Lys Lys Gln

EP 1 382 613 A1

	130	135	140	
5	Lys Ile Leu Asp Ile Arg Lys Asn Val Lys Asp Ala Ile Val Thr Ile			
	145	150	155	160
10	Val Ser Ala Met Ser Thr Ile Ile Pro Pro Val Pro Leu Ala Asn Pro			
	165	170	175	
15	Glu Asn Gln Phe Arg Ser Asp Tyr Ile Lys Ser Ile Ala Pro Ile Thr			
	180	185	190	
20	Asp Phe Glu Tyr Ser Gln Glu Phe Phe Asp His Val Lys Lys Leu Trp			
	195	200	205	
25	Asp Asp Glu Gly Val Lys Ala Cys Phe Glu Arg Ser Asn Glu Tyr Gln			
	210	215	220	
30	Leu Ile Asp Cys Ala Gln Tyr Phe Leu Glu Arg Ile Asp Ser Val Ser			
	225	230	235	240
35	Leu Val Asp Tyr Thr Pro Thr Asp Gln Asp Leu Leu Arg Cys Arg Val			
	245	250	255	
40	Leu Thr Ser Gly Ile Phe Glu Thr Arg Phe Gln Val Asp Lys Val Asn			
	260	265	270	
45	Phe His Met Phe Asp Val Gly Gly Gln Arg Asp Glu Arg Arg Lys Trp			
	275	280	285	
50	Ile Gln Cys Phe Asn Asp Val Thr Ala Ile Ile Tyr Val Ala Ala Cys			
	290	295	300	
55	Ser Ser Tyr Asn Met Val Ile Arg Glu Asp Asn Asn Thr Asn Arg Leu			
	305	310	315	320
	Arg Glu Ser Leu Asp Leu Phe Glu Ser Ile Trp Asn Asn Arg Trp Leu			
	325	330	335	

EP 1 382 613 A1

Arg Thr Ile Ser Ile Ile Leu Phe Leu Asn Lys Gln Asp Met Leu Ala

5 340 345 350

Glu Lys Val Leu Ala Gly Lys Ser Lys Ile Glu Asp Tyr Phe Pro Glu

10 355 360 365

Tyr Ala Asn Tyr Thr Val Pro Glu Asp Ala Thr Pro Asp Ala Gly Glu

15 370 375 380

Asp Pro Lys Val Thr Arg Ala Lys Phe Phe Ile Arg Asp Leu Phe Leu

385 390 395 400

20 Arg Ile Ser Thr Ala Thr Gly Asp Gly Lys His Tyr Cys Tyr Pro His

405 410 415

25 Phe Thr Cys Ala Val Asp Thr Glu Asn Ile Arg Arg Val Phe Asn Asp

420 425 430

30 Cys Arg Asp Ile Ile Gln Arg Met His Leu Lys Gln Tyr Glu Leu Leu

435 440 445

35 <210> 26

<211> 450

40 <212> PRT

<213> Rattus norvegicus

45 <400> 26

Met Gly Leu Cys Tyr Ser Leu Arg Pro Leu Leu Phe Gly Ser Ser Gly

50 1 5 10 15

Asp Ala Pro Cys Glu Asp Ser Glu Pro Cys Ala Glu Asp Ala Gln Pro

55 20 25 30



EP 1 382 613 A1

Ser Ala Ala Pro Ala Pro Ala Pro Ala Pro Ile Pro Ala Pro Ala Pro  
5 35 40 45  
Val Gly Thr Leu Leu Arg Arg Gly Asp Gly Arg Ile Pro Ala Ser Ala  
10 50 55 60  
Arg Ser Pro Val Glu Leu Gln Asn Arg Arg Arg Gln Glu Gln Leu Arg  
15 65 70 75 80  
Ala Glu Glu Arg Glu Ala Ala Lys Glu Ala Arg Lys Val Ser Arg Gly  
85 90 95  
20 Ile Asp Arg Met Leu Arg Glu Gln Lys Arg Asp Leu Gln Gln Thr His  
100 105 110  
25 Arg Leu Leu Leu Leu Gly Ala Gly Glu Ser Gly Lys Ser Thr Ile Val  
115 120 125  
Lys Gln Met Arg Ile Leu His Val Asn Gly Phe Asn Pro Glu Glu Lys  
30 130 135 140  
Lys Gln Lys Ile Leu Asp Ile Arg Lys Asn Val Lys Asp Ala Leu Val  
35 145 150 155 160  
Thr Ile Ile Ser Ala Met Ser Thr Ile Ile Pro Pro Val Pro Leu Ala  
40 165 170 175  
Asn Pro Glu Asn Gln Phe Arg Ser Asp Tyr Ile Lys Ser Ile Ala Pro  
180 185 190  
45 Ile Thr Asp Phe Glu Tyr Ser Gln Glu Phe Phe Asp His Val Lys Lys  
195 200 205  
50 Leu Trp Asp Asp Glu Gly Val Lys Ala Cys Phe Glu Arg Ser Asn Glu  
210 215 220  
55 Tyr Gln Leu Ile Asp Cys Ala Gln Tyr Phe Leu Glu Arg Ile Asp Ser

EP 1 382 613 A1

	225	230	235	240
5	Val Ser Leu Val Asp Tyr Thr Pro Thr Asp Gln Asp Leu Leu Arg Cys			
	245	250	255	
10	Arg Val Leu Thr Ser Gly Ile Phe Glu Thr Arg Phe Gln Val Asp Lys			
	260	265	270	
15	Val Asn Phe His Met Phe Asp Val Gly Gly Gln Arg Asp Glu Arg Arg			
	275	280	285	
20	Lys Trp Ile Gln Cys Phe Asn Asp Val Thr Ala Ile Ile Tyr Val Ala			
	290	295	300	
25	Ala Cys Ser Ser Tyr Asn Met Val Ile Arg Glu Asp Asn Asn Thr Asn			
	305	310	315	320
30	Arg Leu Arg Glu Ser Leu Asp Leu Phe Glu Ser Ile Trp Asn Asn Arg			
	325	330	335	
35	Trp Leu Arg Thr Ile Ser Ile Ile Leu Phe Leu Asn Lys Gln Asp Met			
	340	345	350	
40	Leu Ala Glu Lys Val Leu Ala Gly Lys Ser Lys Ile Glu Asp Tyr Phe			
	355	360	365	
45	Pro Glu Tyr Ala Asn Tyr Thr Val Pro Glu Asp Ala Thr Pro Asp Ala			
	370	375	380	
50	Gly Glu Asp Pro Lys Val Thr Arg Ala Lys Phe Phe Ile Arg Asp Leu			
	385	390	395	400
55	Phe Leu Arg Ile Ser Thr Ala Thr Gly Asp Gly Lys His Tyr Cys Tyr			
	405	410	415	
	Pro His Phe Thr Cys Ala Val Asp Thr Glu Asn Ile Arg Arg Val Phe			
	420	425	430	

Asn Asp Cys Arg Asp Ile Ile Gln Arg Met His Leu Lys Gln Tyr Glu

5 435 440 445

Leu Leu

10 450

<210> 27

15 <211> 1347

<212> DNA

20 <213> Mus musculus

25 <220>

<221> CDS

30 <222> (1).. (1347)

<400> 27

35 atg ggc cta tgc tac agc ctg cgg ccg ctg ctc ttc ggg agc cca gag 48

Met Gly Leu Cys Tyr Ser Leu Arg Pro Leu Leu Phe Gly Ser Pro Glu

40 1 5 10 15

45 gac acc ccg tgt gcg gcc tcg gaa ccc tgc gca gag gat gct cag ccc 96

Asp Thr Pro Cys Ala Ala Ser Glu Pro Cys Ala Glu Asp Ala Gln Pro

50 20 25 30

agc gcc gcc ccg gcc cct gcc tcg atc cca gcc ccg gct ccc gta ggg 144

55 Ser Ala Ala Pro Ala Pro Ala Ser Ile Pro Ala Pro Ala Pro Val Gly

EP 1 382 613 A1

	35	40	45	
5				
	acc ctg ctc cgg cgt ggc ggc ggc cgg atc gtc gcg aac gcg cgg ccg 192			
10	Thr Leu Leu Arg Arg Gly Gly Gly Arg Ile Val Ala Asn Ala Arg Pro			
	50	55	60	
15				
	cca ggc gag ctg cag agc cgc cgg cga cag gag cag cta cga gcc gag 240			
	Pro Gly Glu Leu Gln Ser Arg Arg Arg Gln Glu Gln Leu Arg Ala Glu			
20	65	70	75	80
25				
	gag cgc gag gcg gct aaa gag gcg agg aaa gtc agc cgg ggc atc gac 288			
	Glu Arg Glu Ala Ala Lys Glu Ala Arg Lys Val Ser Arg Gly Ile Asp			
30	85	90	95	
35				
	cgc atg ctg cgc gag cag aag cgg gac ctg cag cag acg cac cgg ctc 336			
	Arg Met Leu Arg Glu Gln Lys Arg Asp Leu Gln Gln Thr His Arg Leu			
40	100	105	110	
45				
	ctg ctg ctg ggg gct ggt gag tcc ggg aaa agc act atc gtc aaa cag 384			
	Leu Leu Leu Gly Ala Gly Glu Ser Gly Lys Ser Thr Ile Val Lys Gln			
50	115	120	125	
55				
	atg agg atc ctg cac gtc aat ggc ttc aac ccc gag gaa aag aag cag 432			
	Met Arg Ile Leu His Val Asn Gly Phe Asn Pro Glu Glu Lys Lys Gln			
	130	135	140	

EP 1 382 613 A1

5       aaa att ctg gac atc agg aaa aat gtc aaa gat gcg atc gtg aca atc   480  
       Lys Ile Leu Asp Ile Arg Lys Asn Val Lys Asp Ala Ile Val Thr Ile  
 10       145                   150                   155                   160

15       gtt tca gca atg agt act atc ata cct cca gtt cca ctg gcc aac cct   528  
       Val Ser Ala Met Ser Thr Ile Ile Pro Pro Val Pro Leu Ala Asn Pro  
                   165                   170                   175

20

25       gag aac cag ttc cgg tca gat tat atc aag agc ata gcc cct atc act   576  
       Glu Asn Gln Phe Arg Ser Asp Tyr Ile Lys Ser Ile Ala Pro Ile Thr  
                   180                   185                   190

30

35       gac ttt gaa tat tcc cag gag ttc ttt gac cat gtg aag aag ctg tgg   624  
       Asp Phe Glu Tyr Ser Gln Glu Phe Phe Asp His Val Lys Lys Leu Trp  
                   195                   200                   205

40       gac gat gaa gga gtg aag gcc tgc ttt gag aga tcc aac gag tac cag   672  
       Asp Asp Glu Gly Val Lys Ala Cys Phe Glu Arg Ser Asn Glu Tyr Gln  
                   210                   215                   220

45

50       ctg atc gac tgt gca caa tac ttc ctg gaa agg att gac agt gtc agt   720  
       Leu Ile Asp Cys Ala Gln Tyr Phe Leu Glu Arg Ile Asp Ser Val Ser  
       225                   230                   235                   240

55

EP 1 382 613 A1

5      ctg gtt gac tac aca ccc aca gac cag gac ctg ctc aga tgc aga gtg    768  
       Leu Val Asp Tyr Thr Pro Thr Asp Gln Asp Leu Leu Arg Cys Arg Val  
                          245                           250                           255

10      ctg aca tca gga atc ttt gag aca cga ttc caa gtg gac aaa gtg aac    816  
       Leu Thr Ser Gly Ile Phe Glu Thr Arg Phe Gln Val Asp Lys Val Asn  
                          260                           265                           270

20      ttt cac atg ttt gat gtt gga ggc cag aga gat gag aga aga aaa tgg    864  
       Phe His Met Phe Asp Val Gly Gly Gln Arg Asp Glu Arg Arg Lys Trp  
                          275                           280                           285

30      atc cag tgt ttt aat gat gtc act gcg atc att tac gtg gcg gcc tgt    912  
       Ile Gln Cys Phe Asn Asp Val Thr Ala Ile Ile Tyr Val Ala Ala Cys  
                          290                           295                           300

40      agt agc tac aac atg gtg atc cgg gaa gat aac aat acc aac aga ctt    960  
       Ser Ser Tyr Asn Met Val Ile Arg Glu Asp Asn Asn Thr Asn Arg Leu  
                          305                           310                           315                           320

50      cgg gaa tca ctg gac ctg ttt gaa agc atc tgg aat aac agg tgg ttg    1008  
       Arg Glu Ser Leu Asp Leu Phe Glu Ser Ile Trp Asn Asn Arg Trp Leu  
                          325                           330                           335

55      cga acc att tct atc atc cta ttc ttg aac aaa caa gac atg ctg gca    1056

EP 1 382 613 A1

Arg Thr Ile Ser Ile Ile Leu Phe Leu Asn Lys Gln Asp Met Leu Ala

5 340 345 350

10 gaa aaa gtc ttg gca ggg aag tca aaa atc gaa gac tat ttc ccg gag 1104

Glu Lys Val Leu Ala Gly Lys Ser Lys Ile Glu Asp Tyr Phe Pro Glu

15 355 360 365

20 tat gcc aat tat act gtc cct gaa gat gca aca cca gat gcg gga gaa 1152

Tyr Ala Asn Tyr Thr Val Pro Glu Asp Ala Thr Pro Asp Ala Gly Glu

25 370 375 380

gat ccc aaa gtt aca aga gca aag ttc ttt atc cgg gat ctg ttc ttg 1200

30 Asp Pro Lys Val Thr Arg Ala Lys Phe Phe Ile Arg Asp Leu Phe Leu

385 390 395 400

35 agg atc agc aca gcc acg ggt gat ggc aaa cat tac tgc tac cct cac 1248

Arg Ile Ser Thr Ala Thr Gly Asp Gly Lys His Tyr Cys Tyr Pro His

40 405 410 415

45 ttc acc tgc gcc gtg gac aca gag aac atc cgc aga gtg ttc aac gat 1296

Phe Thr Cys Ala Val Asp Thr Glu Asn Ile Arg Arg Val Phe Asn Asp

50 420 425 430

55 tgc cgt gac atc atc cag aga atg cat ctc aag cag tac gaa ctc ttg 1344

Cys Arg Asp Ile Ile Gln Arg Met His Leu Lys Gln Tyr Glu Leu Leu

435

440

445

tga

1347

&lt;210&gt; 28

&lt;211&gt; 1353

&lt;212&gt; DNA

&lt;213&gt; Rattus norvegicus

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (1).. (1353)

&lt;400&gt; 28

atg ggc ctg tgc tac agc cta cgg ccg ctg ctc ttc ggg agc tcg ggg 48

Met Gly Leu Cys Tyr Ser Leu Arg Pro Leu Leu Phe Gly Ser Ser Gly

1 5 10 15

gac gcc ccc tgt gag gac tct gag ccg tgc gct gag gat gct cag ccc 96

Asp Ala Pro Cys Glu Asp Ser Glu Pro Cys Ala Glu Asp Ala Gln Pro

20 25 30

agc gcc gcc ccg gcc ccg gcc ccg gcc ccg atc cca gcc ccg gct ccg 144

Ser Ala Ala Pro Ala Pro Ala Pro Ala Pro Ile Pro Ala Pro Ala Pro

35 40 45



EP 1 382 613 A1

gtg ggg acc ctg ctc egg cga ggc gac ggc egg atc ccc gca agc gcg 192  
5 Val Gly Thr Leu Leu Arg Arg Gly Asp Gly Arg Ile Pro Ala Ser Ala  
50 55 60

agg tcg cca gtc gag ctg cag aac cgc egg cga cag gag cag ctg cga 240  
10 Arg Ser Pro Val Glu Leu Gln Asn Arg Arg Arg Gln Glu Gln Leu Arg  
15 65 70 75 80

gcc gag gag cgc gag gca gct aag gag gcg agg aaa gta agc egg ggt 288  
20 Ala Glu Glu Arg Glu Ala Ala Lys Glu Ala Arg Lys Val Ser Arg Gly  
25 85 90 95

atc gac cgc atg ctg cgc gaa cag aag cgc gac ctg cag cag acg cac 336  
30 Ile Asp Arg Met Leu Arg Glu Gln Lys Arg Asp Leu Gln Gln Thr His  
35 100 105 110

egg ctc ctg ctc ttg ggg gct ggt gag tcc ggg aaa agc act ata gtc 384  
40 Arg Leu Leu Leu Leu Gly Ala Gly Glu Ser Gly Lys Ser Thr Ile Val  
45 115 120 125

aaa cag atg agg atc cta cac gtc aat ggc ttc aac ccc gag gaa aag 432  
50 Lys Gln Met Arg Ile Leu His Val Asn Gly Phe Asn Pro Glu Glu Lys  
55 130 135 140

EP 1 382 613 A1

aag cag aaa att ctg gac atc agg aaa aat gtc aaa gat gct tta gtg 480  
 5 Lys Gln Lys Ile Leu Asp Ile Arg Lys Asn Val Lys Asp Ala Leu Val  
 145 150 155 160  
 10 aca atc att tca gca atg agt acc ata ata cct cca gtt cca ctg gcc 528  
 Thr Ile Ile Ser Ala Met Ser Thr Ile Ile Pro Pro Val Pro Leu Ala  
 15 165 170 175  
 20 aac cct gag aac cag ttt cgg tca gat tac atc aag agc ata gcc cct 576  
 Asn Pro Glu Asn Gln Phe Arg Ser Asp Tyr Ile Lys Ser Ile Ala Pro  
 25 180 185 190  
 30 atc act gac ttt gaa tat tcc cag gag ttc ttt gac cac gtg aag aag 624  
 Ile Thr Asp Phe Glu Tyr Ser Gln Glu Phe Phe Asp His Val Lys Lys  
 195 200 205  
 35 ctg tgg gat gat gag gga gtg aag gcc tgc ttt gag aga tcc aac gag 672  
 40 Leu Trp Asp Asp Glu Gly Val Lys Ala Cys Phe Glu Arg Ser Asn Glu  
 210 215 220  
 45 tac cag ctg atc gac tgt gca caa tac ttc ctg gaa agg att gac agc 720  
 Tyr Gln Leu Ile Asp Cys Ala Gln Tyr Phe Leu Glu Arg Ile Asp Ser  
 50 225 230 235 240  
 55 gtg agt ctg gtt gac tac aca ccc aca gac cag gac cta ctc aga tgc 768

EP 1 382 613 A1

	Val Ser Leu Val Asp Tyr Thr Pro Thr Asp Gln Asp Leu Leu Arg Cys	
5	245	250 255
10	aga gtg ctg aca tca ggg atc ttt gag aca cga ttc caa gtg gac aaa	816
	Arg Val Leu Thr Ser Gly Ile Phe Glu Thr Arg Phe Gln Val Asp Lys	
15	260	265 270
20	gtg aac ttt cac atg ttt gac gtt gga ggc cag agg gat gag aga aga	864
	Val Asn Phe His Met Phe Asp Val Gly Gly Gln Arg Asp Glu Arg Arg	
25	275	280 285
30	aaa tgg atc cag tgt ttt aac gat gtc act gcc atc atc tat gtg gca	912
	Lys Trp Ile Gln Cys Phe Asn Asp Val Thr Ala Ile Ile Tyr Val Ala	
35	290	295 300
40	gcc tgc agc agc tac aac atg gtg atc cgg gaa gat aac aac acc aac	960
	Ala Cys Ser Ser Tyr Asn Met Val Ile Arg Glu Asp Asn Asn Thr Asn	
45	305	310 315 320
50	aga ctc cgg gag tgc ctg gac ctg ttt gaa agc atc tgg aat aac agg	1008
	Arg Leu Arg Glu Ser Leu Asp Leu Phe Glu Ser Ile Trp Asn Asn Arg	
55	325	330 335
	tgg tta cga acc att tcc atc atc ctg ttc ttg aac aaa caa gat atg	1056
	Trp Leu Arg Thr Ile Ser Ile Ile Leu Phe Leu Asn Lys Gln Asp Met	

EP 1 382 613 A1

	340	345	350	
5				
	ctg gca gaa aaa gtc ttg gcc ggg aag tca aaa att gaa gac tat ttc	1104		
10	Leu Ala Glu Lys Val Leu Ala Gly Lys Ser Lys Ile Glu Asp Tyr Phe			
	355	360	365	
15				
	cag gag tat gcc aac tat act gtc cct gaa gat gca aca cca gat gca	1152		
	Pro Glu Tyr Ala Asn Tyr Thr Val Pro Glu Asp Ala Thr Pro Asp Ala			
20	370	375	380	
25				
	gga gaa gat ccc aaa gtt aca aga gcc aag ttc ttt atc cgg gat ctg	1200		
	Gly Glu Asp Pro Lys Val Thr Arg Ala Lys Phe Phe Ile Arg Asp Leu			
30	385	390	395	400
35				
	ttc ttg agg atc agc aca gcc acg ggt gat ggc aaa cat tac tgc tac	1248		
	Phe Leu Arg Ile Ser Thr Ala Thr Gly Asp Gly Lys His Tyr Cys Tyr			
40	405	410	415	
45				
	cct cac ttc acc tgc gcc gtg gac aca gag aac atc cgc aga gtg ttc	1296		
	Pro His Phe Thr Cys Ala Val Asp Thr Glu Asn Ile Arg Arg Val Phe			
	420	425	430	
50				
	aac gat tgt cgt gac atc atc cag aga atg cac ctc aag cag tac gaa	1344		
	Asn Asp Cys Arg Asp Ile Ile Gln Arg Met His Leu Lys Gln Tyr Glu			
55	435	440	445	

ctc ttg tga

1353

5

Leu Leu

450

10

<210> 29

15

<211> 29

<212> DNA

20

<213> Artificial Sequence

<220>

25

<223> a primer used in an example of the present invention

30

<400> 29

atgggcctat gctacagcct gcggccgct

29

35

<210> 30

<211> 33

40

<212> DNA

<213> Artificial Sequence

45

<220>

50

<223> a primer used in an example of the present invention

55

<400> 30

tcacaagagt tcgtactgct tgagatgcat tct

33

5

<210> 31

10

<211> 30

<212> DNA

15

<213> Artificial Sequence

<220>

20

<223> a primer used in an example of the present invention

25

<400> 31

atgggcctgt gctacagcct acggccgctg

30

30

<210> 32

<211> 33

35

<212> DNA

<213> Artificial Sequence

40

<220>

45

<223> a primer used in an example of the present invention

50

<400> 32

tcacaagagt tcgtactgct tgaggatgcat tct

33

55

<210> 33

<211> 37

<212> DNA

<213> Artificial Sequence

<220>

<223> a primer used in an example of the present invention

<400> 33

agctcggatc catgcccatc atgggctcct cgggtga

37

<210> 34

<211> 34

<212> DNA

<213> Artificial Sequence

<220>

<223> a primer used in an example of the present invention

<400> 34

gtgcagaatt ctcaggacac tctgctcca tcct

34

# Claims

1. A protein comprising any amino acid sequence selected from the group consisting of:

- (a) the amino acid sequence represented by SEQ ID No:1;
- (b) an amino acid sequence of a protein involved in a G protein-coupled receptor mediated signal transduction, said protein consists of an amino acid sequence having a homology of 85% or more with the amino acid sequence represented by SEQ ID No:1;
- (c) the amino acid sequence represented by SEQ ID No:25;

- (d) the amino acid sequence represented by SEQ ID No:26;
- (e) an amino acid sequence of a protein involved in a G protein-coupled receptor mediated signal transduction, said protein comprises the amino acid sequence of the amino acid Nos.96 to 126 of SEQ ID No:1;
- (f) an amino acid sequence of a protein involved in a G protein-coupled receptor mediated signal transduction, said protein comprises an amino acid sequence having a homology of 95% or more with the amino acid sequence of the amino acid Nos.96 to 126 of SEQ ID No:1;
- (g) an amino acid sequence of a protein involved in a G protein-coupled receptor mediated signal transduction, said protein comprises at its N-terminal the amino acid sequence of the amino acid Nos.1 to 126 of SEQ ID No:1; and
- (h) an amino acid sequence of a protein involved in a G protein-coupled receptor mediated signal transduction, said protein comprises at its N-terminal an amino acid sequence having a homology of 65% or more with the amino acid sequence of the amino acid Nos.1 to 126 of SEQ ID No:1.

2. A protein (1) or (2):

- (1) the protein consisting of the amino acid sequence represented by SEQ ID No:1;
- (2) a protein involved in a G protein-coupled receptor mediated signal transduction which consists of an amino acid sequence having a homology of 85% or more with the amino acid sequence represented by SEQ ID No:1.

3. The protein consisting of the amino acid sequence represented by SEQ ID No:25.

4. The protein consisting of the amino acid sequence represented by SEQ ID No:26.

5. A protein (3) or (4):

- (3) a protein involved in a G protein-coupled receptor mediated signal transduction which comprises the amino acid sequence of the amino acid Nos.96 to 126 of SEQ ID No:1;
- (4) a protein involved in a G protein-coupled receptor mediated signal transduction which comprises an amino acid sequence having a homology of 95% or more with the amino acid sequence of the amino acid Nos.96 to 126 of SEQ ID No:1.

6. A protein (5) or (6):

- (5) a protein involved in a G protein-coupled receptor mediated signal transduction which comprises at its N-terminal the amino acid sequence of the amino acid Nos.1 to 126 of SEQ ID No:1;
- (6) a protein involved in a G protein-coupled receptor mediated signal transduction which comprises at its N-terminal an amino acid sequence having a homology of 65% or more with the amino acid sequence of the amino acid Nos.1 to 126 of SEQ ID No:1.

7. A polynucleotide comprising a nucleotide sequence selected from the group consisting of:

- (a) a nucleotide sequence encoding the amino acid sequence represented by SEQ ID No:1;
- (b) a nucleotide sequence encoding an amino acid sequence of a protein involved in a G protein-coupled receptor mediated signal transduction, said protein consists of an amino acid sequence having a homology of 85% or more with the amino acid sequence represented by SEQ ID No:1;
- (c) a nucleotide sequence encoding the amino acid sequence represented by SEQ ID No:25;
- (d) a nucleotide sequence encoding the amino acid sequence represented by SEQ ID No:26;
- (e) a nucleotide sequence encoding an amino acid sequence of a protein involved in a G protein-coupled receptor mediated signal transduction, said protein comprises the amino acid sequence of the amino acid Nos.96 to 126 of SEQ ID No:1;
- (f) a nucleotide sequence encoding an amino acid sequence of a protein involved in a G protein-coupled receptor mediated signal transduction, said protein comprises an amino acid sequence having a homology of 95% or more with the amino acid sequence of the amino acid Nos.96 to 126 of SEQ ID No:1;
- (g) a nucleotide sequence encoding an amino acid sequence of a protein involved in a G protein-coupled receptor mediated signal transduction, said protein comprises at its N-terminal the amino acid sequence of the amino acid Nos.1 to 126 of SEQ ID No:1;
- (h) a nucleotide sequence encoding an amino acid sequence of a protein involved in a G protein-coupled receptor mediated signal transduction, said protein comprises at its N-terminal an amino acid sequence having



a homology of 65% or more with the amino acid sequence of the amino acid Nos.1 to 126 of SEQ ID No:1;  
 (i) the nucleotide sequence represented by SEQ ID No:2;  
 (j) a nucleotide sequence of a polynucleotide encoding a protein involved in a G protein-coupled receptor mediated signal transduction, said polynucleotide consists of a nucleotide sequence having a homology of 85% or more with the polynucleotide consisting of the nucleotide sequence represented by SEQ ID No:2;  
 (k) the nucleotide sequence represented by SEQ ID No:27;  
 (l) the nucleotide sequence represented by SEQ ID No:28;  
 (m) a nucleotide sequence of a polynucleotide encoding a protein involved in a G protein-coupled receptor mediated signal transduction, said polynucleotide comprises the nucleotide sequence of the nucleotide Nos. 289 to 378 of SEQ ID No:2;  
 (n) a nucleotide sequence of a polynucleotide encoding a protein involved in a G protein-coupled receptor mediated signal transduction, said polynucleotide comprises a nucleotide sequence having a homology of 90% or more with the polynucleotide consisting of the nucleotide sequence of the nucleotide Nos.289 to 378 of SEQ ID No:2;  
 (o) a nucleotide sequence of a polynucleotide encoding a protein involved in a G protein-coupled receptor mediated signal transduction, said polynucleotide comprises at its 5' terminal the nucleotide sequence of the nucleotide Nos.1 to 378 of SEQ ID No:2; and  
 (p) a nucleotide sequence of a polynucleotide encoding a protein involved in a G protein-coupled receptor mediated signal transduction, said polynucleotide comprises at its 5' terminal a nucleotide sequence having a homology of 70% or more with the polynucleotide consisting of the nucleotide sequence of the nucleotide Nos.1 to 378 of SEQ ID No:2.

8. A polynucleotide of (7) or (8):

(7) the polynucleotide consisting of the nucleotide sequence represented by SEQ ID No:2;  
 (8) a polynucleotide encoding a protein involved in a G protein-coupled receptor mediated signal transduction, said polynucleotide consists of a nucleotide sequence having a homology of 85% or more with the polynucleotide consisting of the nucleotide sequence represented by SEQ ID No:2.

9. The polynucleotide consisting of the nucleotide sequence represented by SEQ ID No:27.

10. The polynucleotide consisting of the nucleotide sequence represented by SEQ ID No:28.

11. A polynucleotide of (9) or (10):

(9) a polynucleotide encoding a protein involved in a G protein-coupled receptor mediated signal transduction, said polynucleotide comprises the nucleotide sequence of the nucleotide Nos.289 to 378 of SEQ ID No:2;  
 (10) a polynucleotide encoding a protein involved in a G protein-coupled receptor mediated signal transduction, said polynucleotide comprises a nucleotide sequence having a homology of 90% or more with the polynucleotide consisting of the nucleotide sequence of the nucleotide Nos.289 to 378 of SEQ ID No:2.

12. A polynucleotide of (11) or (12):

(11) a polynucleotide encoding a protein involved in a G protein-coupled receptor mediated signal transduction, said polynucleotide comprises at its 5' terminal the nucleotide sequence of the nucleotide Nos.1 to 378 of SEQ ID No:2;  
 (12) a polynucleotide encoding a protein involved in a G protein-coupled receptor mediated signal transduction, said polynucleotide comprises at its 5' terminal a nucleotide sequence having a homology of 70% or more with the polynucleotide consisting of the nucleotide sequence of the nucleotide Nos.1 to 378 of SEQ ID No:2.

13. A recombinant vector containing a polynucleotide according to Claim 7.

14. A method for producing a recombinant vector comprising a step for integrating a polynucleotide according to Claim 7 into a vector capable of being replicated in a host cell.

15. A transformant having a recombinant vector according to claim 13.

16. A method for producing a transformant comprising a step for transducing a recombinant vector according to claim

13 into a host cell.

17. A method for producing a G protein  $\alpha$ -subunit comprising steps for culturing a transformant having a recombinant vector containing a polynucleotide according to Claim 7 and recovering from the culture a protein resulting from the polynucleotide according to Claim 7.

18. An antisense polynucleotide consisting of a polynucleotide of (13) or (14):

(13) a polynucleotide which inhibits the expression of a protein according to Claim 1 which comprises a nucleotide sequence complementary to at least 15 contiguous nucleotides in the nucleotide sequence represented by SEQ ID No:2;

(14) a polynucleotide which inhibits the expression of a protein according to Claim 1 which hybridizes under an intracellular condition with a polynucleotide consisting of at least 15 contiguous nucleotides in the nucleotide sequence represented by SEQ ID No:2.

19. A ribozyme (15) or (16):

(15) a ribozyme having an ability of cleaving a polynucleotide according to Claim 7 which comprises two polynucleotide regions complementary to two regions respectively consisting of at least 9 contiguous nucleotides which are two regions in the nucleotide sequence represented by SEQ ID No:2;

(16) a ribozyme having an ability of cleaving a polynucleotide according to Claim 7 which comprises two polynucleotide regions which hybridizes under an intracellular condition with two regions respectively consisting of at least 9 contiguous nucleotides which are two regions in the nucleotide sequence represented by SEQ ID No:2.

20. An antibody which recognizes a protein according to Claim 1 specifically.

21. An agent for regulating a G protein-coupled receptor mediated signal transduction containing as an active ingredient a protein according to Claim 1.

22. A therapeutic or prophylactic agent against a disease caused by a G protein-coupled receptor mediated signal transduction abnormality, wherein an active ingredient of the agent is a protein according to Claim 1.

23. An agent for regulating a G protein-coupled receptor mediated signal transduction containing as an active ingredient a polynucleotide according to Claim 7.

24. A therapeutic or prophylactic agent against a disease caused by a G protein-coupled receptor mediated signal transduction abnormality, wherein an active ingredient of the agent is a polynucleotide according to Claim 7.

25. An agent for regulating a G protein-coupled receptor mediated signal transduction containing as an active ingredient an antisense polynucleotide according to Claim 18.

26. A therapeutic or prophylactic agent against a disease caused by a G protein-coupled receptor mediated signal transduction abnormality, wherein an active ingredient of the agent is an antisense polynucleotide according to Claim 18.

27. An agent for regulating a G protein-coupled receptor mediated signal transduction containing as an active ingredient a ribozyme according to Claim 19.

28. A therapeutic or prophylactic agent against a disease caused by a G protein-coupled receptor mediated signal transduction abnormality, wherein an active ingredient of the agent is a ribozyme according to Claim 19.

29. An agent for regulating a G protein-coupled receptor mediated signal transduction containing as an active ingredient an antibody according to Claim 20.

30. A therapeutic or prophylactic agent against a disease caused by a G protein-coupled receptor mediated signal transduction abnormality, wherein an active ingredient of the agent is an antibody according to Claim 20.

31. An oligonucleotide (17) or (18):

(17) an oligonucleotide capable of recognizing a polynucleotide represented by SEQ ID NO:2 specifically which consists of at least 17 contiguous nucleotides in the nucleotide sequence represented by SEQ ID No:2;  
 (18) an oligonucleotide capable of recognizing a polynucleotide represented by SEQ ID NO:2 specifically which has a homology of 80% or more with at least 17 contiguous nucleotides in the nucleotide sequence represented by SEQ ID No:2.

32. An oligonucleotide according to Claim 31 which is used as a probe or a primer.

33. A method for screening for a substance capable of regulating a signal transduction mediated by a G protein-coupled receptor and a G protein comprising:

(a) a step for bringing a test substance into contact with a test cell having a recombinant vector according to Claim 13 and a recombinant vector containing a DNA encoding a G protein-coupled receptor protein;  
 (b) a step for measuring the G protein effector activity or the index value correlating therewith in the test cell; and  
 (c) a step for comparing this effector activity or the index value correlating therewith with the effector activity or the index value correlating therewith in the test cell which has not been brought into contact with the test substance, whereby selecting a test substance capable of altering the effector activity or the index value correlating therewith in the test cell.

34. A method for screening for a substance capable of regulating a signal transduction mediated by a G protein-coupled receptor and a G protein comprising:

(a) a step for bringing a test substance into contact with a test cell having a recombinant vector according to Claim 13 and a recombinant vector containing a DNA encoding a G protein-coupled receptor protein;  
 (b) a step for measuring the G protein effector activity or the index value correlating therewith in the test cell; and  
 (c) a step for comparing this effector activity with the effector activity or the index value correlating therewith when the said test substance has been brought into contact with a control cell having no recombinant vector according to Claim 13 but having a recombinant vector containing a DNA encoding a G protein-coupled receptor protein, whereby selecting a test substance causing a difference in the effector activity or the index value correlating therewith between the test cell and the control cell.

35. A method for screening for a substance capable of regulating a signal transduction mediated by a G protein-coupled receptor and a G protein comprising:

(a) a step for bringing a test substance into contact with a test cell having a recombinant vector according to Claim 13 and a recombinant vector containing a DNA encoding a G protein-coupled receptor protein;  
 (b) a step for measuring the G protein effector activity or the index value correlating therewith in the test cell; and  
 (c) a step for comparing this effector activity or the index value correlating therewith with the effector activity or the index value correlating therewith when the said test substance has been brought into contact with a control cell having no recombinant vector containing a DNA encoding a G protein-coupled receptor protein but having a recombinant vector according to Claim 13, whereby selecting a test substance causing a difference in the effector activity or the index value correlating therewith between the test cell and the control cell.

36. A method for screening for a substance capable of regulating a signal transduction mediated by a G protein-coupled receptor and a G protein comprising:

(a) a step for bringing a test substance and a G protein-coupled receptor ligand into contact with a test cell having a recombinant vector according to Claim 13 and a recombinant vector containing a DNA encoding a G protein-coupled receptor protein;  
 (b) a step for measuring the G protein effector activity or the index value correlating therewith in the test cell; and  
 (c) a step for comparing this effector activity or the index value correlating therewith with the effector activity or the index value correlating therewith in the test cell which has not been brought into contact with the test substance but has been brought into contact with the ligand, whereby selecting a test substance capable of altering the effector activity or the index value correlating therewith in the test cell.

37. A method for screening for a substance capable of regulating a signal transduction mediated by a G protein-coupled

receptor and a G protein comprising:

- (a) a step for bringing a test substance and a G protein-coupled receptor ligand into contact with a test cell having a recombinant vector according to Claim 13 and a recombinant vector containing a DNA encoding a G protein-coupled receptor protein;
- (b) a step for measuring the G protein effector activity or the index value correlating therewith in the test cell;
- (c) a step for comparing this effector activity with the effector activity in the test cell which has not been brought into contact with the test substance but has been brought into contact with the ligand, whereby investigating the change in the effector activity in the test cell; and
- (d) a step for comparing the rate of change in this effector activity or the index value correlating therewith with the rate of change in the effector activity or the index value correlating therewith when the said test substance and said ligand has been brought into contact with a control cell having no recombinant vector containing a DNA encoding a G protein-coupled receptor protein but having a recombinant vector according to Claim 13, whereby selecting a test substance causing a difference in the rate of change in the effector activity or the index value correlating therewith between the test cell and the control cell.

38. A method for screening for a substance capable of regulating a signal transduction mediated by a G protein-coupled receptor and a G protein comprising:

- (a) a step for bringing a test substance into contact with a cell membrane fraction of a cell having a recombinant vector according to Claim 13 and a cell membrane fraction of a cell having a recombinant vector containing a DNA encoding a GPCR, or a cell membrane fraction of a cell having the recombinant vector according to Claim 13 and the recombinant vector containing the DNA encoding the GPCR;
- (b) a step for assaying the level of the binding of GTP to the cell membrane fraction; and
- (c) a step for comparing the assayed level of this GTP binding with the assayed level of the GTP binding to the cell membrane fraction which has not been brought into contact with the test substance, whereby selecting a test substance capable of altering the assayed level of the GTP binding to the cell membrane fraction.

39. A method for screening for a substance capable of regulating a signal transduction mediated by a G protein-coupled receptor and a G protein comprising:

- (a) a step for bringing a test substance and a G protein-coupled receptor ligand into contact with a cell membrane fraction of a cell having a recombinant vector according to Claim 13 and a cell membrane fraction of a cell having a recombinant vector containing a DNA encoding a GPCR, or a cell membrane fraction of a cell having the recombinant vector according to Claim 13 and the recombinant vector containing the DNA encoding the GPCR;
- (b) a step for assaying the level of the binding of GTP to the cell membrane fraction; and
- (c) a step for comparing the assayed level of this GTP binding with the assayed level of the GTP binding in the cell membrane fraction which has not been brought into contact with the test substance but has been brought into contact with said ligand, whereby selecting a test substance capable of altering the assayed level of the GTP binding to the cell membrane fraction.

40. A method for screening for a substance capable of regulating a signal transduction mediated by a G protein-coupled receptor and a G protein comprising:

- (a) a step for bringing a test substance into contact with a test cell capable of expressing a protein according to Claim 1;
- (b) a step for measuring the expression level of the protein according to Claim 1 in the test cell; and
- (c) a step for comparing this expression level with the expression level of said protein in the test cell which has not been brought into contact with the test substance, whereby selecting a test substance capable of altering the expression level of said protein in the test cell.

41. A substance capable of regulating a signal transduction mediated by a G protein-coupled receptor and a G protein, said substance is obtained by a screening method according to any of Claims 33 to 40.

42. An agent for regulating a signal transduction mediated by a G protein-coupled receptor and a G protein, said agent contains as an active ingredient a substance according to Claim 41.

43. A therapeutic or prophylactic agent against a disease caused by the abnormality in a G protein-coupled receptor and a G protein-mediated signal transduction containing as an active ingredient a substance according to Claim 41.

44. A kit for screening for a substance capable of regulating a signal transduction mediated by a G protein-coupled receptor and a protein according to Claim 1, which comprises a test cell having a recombinant vector containing a polynucleotide encoding a protein according to Claim 1 and a reagent for measuring the G protein effector activity or an index value correlating therewith.

45. A screening kit according to Claim 44 wherein the test cell further has a recombinant vector having a DNA encoding a G protein-coupled receptor.

46. A screening kit according to Claim 44 further containing a G protein-coupled receptor ligand.

47. A screening kit according to Claim 44 further containing a control cell having a recombinant vector having a DNA encoding a G protein-coupled receptor.

48. A screening kit according to Claim 44 further containing a control cell having a recombinant vector containing a polynucleotide encoding a protein according to Claim 1.

49. A kit for screening for a substance capable of regulating a signal transduction mediated by a G protein-coupled receptor and a protein according to Claim 1, which comprises:

a cell having a recombinant vector containing a polynucleotide encoding a protein according to Claim 1; and, a GTP analogue which can bind to the protein according to Claim 1 but can not be cleaved by a GTPase.

50. A screening kit according to Claim 49 wherein the cell further has a recombinant vector having a DNA encoding a G protein-coupled receptor.

51. A kit for screening for a substance capable of regulating a signal transduction mediated by a G protein-coupled receptor and a protein according to Claim 1, which comprises:

a cell having a recombinant vector containing a polynucleotide encoding a protein according to Claim 1; a cell having a recombinant vector having a DNA encoding the G protein-coupled receptor; and, a GTP analogue which can bind to the protein according to Claim 1 but can not be cleaved by a GTPase.

52. A screening kit according to Claim 49 or 51 further containing a G protein-coupled receptor ligand.

53. A kit for screening for a substance capable of regulating a signal transduction mediated by a G protein-coupled receptor and a protein according to Claim 1, which comprises:

a cell capable of expressing a protein according to Claim 1; and an oligonucleotide according to Claim 31 or an antibody according to Claim 20.

Figure 1




Figure 2




Figure 3




Figure 4




Figure 5




Figure 6







Figure 7





European Patent  
Office

# PARTIAL EUROPEAN SEARCH REPORT

Application Number

which under Rule 45 of the European Patent Convention EP 03 01 5519  
shall be considered, for the purposes of subsequent  
proceedings, as the European search report

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
X	JONES D T ET AL: "GOLF: AN OLFACTORY NEURON SPECIFIC-G PROTEIN INVOLVED IN ODORANT SIGNAL TRANSDUCTION" SCIENCE, AMERICAN ASSOCIATION FOR THE ADVANCEMENT OF SCIENCE,, US, vol. 244, no. 4906, 19 May 1989 (1989-05-19), pages 790-795, XP001109598 ISSN: 0036-8075	18,25, 26,31,32	C07K14/47
Y	* page 2509 - page 2510; figure 1 *	1-53	
X	-& DATABASE EMBL [Online] Human GTP binding protein, retrieved from EBI Database accession no. L10665 XP002260433	18,25, 26,31,32	
			TECHNICAL FIELDS SEARCHED (Int.Cl.7)
			C07K
<b>INCOMPLETE SEARCH</b>			
<p>The Search Division considers that the present application, or one or more of its claims, does/do not comply with the EPO to such an extent that a meaningful search into the state of the art cannot be carried out, or can only be carried out partially, for these claims.</p> <p>Claims searched completely :</p> <p>Claims searched incompletely :</p> <p>Claims not searched :</p> <p>Reason for the limitation of the search:</p> <p>see sheet C</p>			
Place of search		Date of completion of the search	Examiner
MUNICH		6 November 2003	Trommsdorff, M
<p><b>CATEGORY OF CITED DOCUMENTS</b></p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technological background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons</p> <p>&amp; : member of the same patent family, corresponding document</p>			

EPO FORM 1 503 03 82 (P04C37)



European Patent  
Office

**INCOMPLETE SEARCH  
SHEET C**

Application Number

EP 03 01 5519

Claim(s) not searched:  
41-43

Reason for the limitation of the search:

Present claims 41-43 relate to a substances defined by reference to a desirable characteristic or property, namely their ability to regulate a signal transduction mediated by a GPCR and a G protein. The claims cover all substances having this characteristic or property, whereas the application does not provide support within the meaning of Article 84 EPC and/or disclosure within the meaning of Article 83 EPC for any such substances. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search is impossible. Independent of the above reasoning, the claims also lack clarity (Article 84 EPC). An attempt is made to define the substances by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search impossible. Consequently, no search has been carried out for those claims.



European Patent  
Office

## PARTIAL EUROPEAN SEARCH REPORT

Application Number  
EP 03 01 5519

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int.Cl.7)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
X	ZIGMAN J M ET AL: "HUMAN GOLFALPHA: COMPLEMENTARY DEOXYRIBONUCLEIC ACID STRUCTURE AND EXPRESSION IN PANCREATIC ISLETS AND OTHER TISSUES OUTSIDE THE OLFACTORY NEUROEPITHELIUM AND CENTRAL NERVOUS SYSTEM" ENDOCRINOLOGY, BALTIMORE, MD, US, vol. 133, no. 8, December 1993 (1993-12), pages 2508-2514, XP000993079 ISSN: 0013-7227 * page 2509 - page 2510 *	18,25, 26,31,32	
X	WO 02 22882 A (ARCHEMIX CORP ;EPSTEIN DAVID (US); STANTON MARTY (US); HAMAGUCHI N) 21 March 2002 (2002-03-21) * page 88 - page 89; example 12 *	27,28	TECHNICAL FIELDS SEARCHED (Int.Cl.7)
X	US 6 110 664 A (COWSERT LEX M) 29 August 2000 (2000-08-29) * example 15; table 1 *	17,18	
Y	BOURNE H R ET AL: "THE GTPASE SUPERFAMILY CONSERVED STRUCTURE AND MOLECULAR MECHANISM" NATURE (LONDON), vol. 349, no. 6305, 1991, pages 117-127, XP001153219 ISSN: 0028-0836 * the whole document *	1-53	
P,X	STRAUSBERG R L ET AL: "Generation and initial analysis of more than 15,000 full-length human and mouse cDNA sequences" NCBI, XP002245204 * the whole document *	1-53	

EPO FORM 1503 03.02 (P/C10)

**ANNEX TO THE EUROPEAN SEARCH REPORT  
ON EUROPEAN PATENT APPLICATION NO.**

EP 03 01 5519

This annex lists the patent family members relating to the patent documents cited in the above-mentioned European search report. The members are as contained in the European Patent Office EDP file on  
The European Patent Office is in no way liable for these particulars which are merely given for the purpose of information.

06-11-2003

Patent document cited in search report		Publication date		Patent family member(s)	Publication date
WO 0222882	A	21-03-2002	AU	9096501 A	26-03-2002
			CA	2418724 A1	21-03-2002
			EP	1354062 A2	22-10-2003
			WO	0222882 A2	21-03-2002
			US	2003087239 A1	08-05-2003
-----					
US 6110664	A	29-08-2000	AU	5486700 A	31-01-2001
			WO	0100861 A1	04-01-2001
-----					

EPO FORM P0459

For more details about this annex : see Official Journal of the European Patent Office, No. 12/82

